


12

USER MANUAL



PARASURFTM

Impressum

Copyright

© 2013 by Cepos InSilico Ltd.

The Old Vicarage
132 Bedford Road
Kempston
BEDFORD, MK42 8BQ
United Kingdom
www.ceposinsilico.com

Manual

Timothy Clark

Layout

www.eh-bitartist.de





TABLE OF CONTENTS

PROGRAM HISTORY	5
1 INTRODUCTION	6
1.1 Changes relative to ParaSurf'11™	7
1.1.1 Updated local properties	7
1.1.2 Local polarizability: new definition	7
1.1.3 Electron density: scaling error	7
1.1.4 EMPIRE™	7
1.1.5 HDF5 files	7
1.2 Isodensity surfaces	8
1.3 Solvent-excluded surfaces	8
1.4 Solvent-accessible surfaces	8
1.5 Shrink-wrap surface algorithm	9
1.6 Marching-cube algorithm	10
1.7 Spherical-harmonic fitting	11
1.8 Local properties	13
1.8.1 Molecular electrostatic potential	13
1.8.1.1 The natural atomic orbital/PC (NAO-PC) model	13
1.8.1.2 The multipole model	13
1.8.2 Local ionization energy, electron affinity, hardness and electronegativity	14
1.8.3 Local polarizability	15
1.8.4 Field normal to the surface	15
1.9 Descriptors	16
1.10 Surface-integral models (polynomial version)	21
1.11 Binned surface-integral models	22
1.12 Spherical harmonic "hybrids"	23
1.13 Descriptors and moments based on polynomial surface-integral models	24
1.14 Shannon entropy	25
1.15 Surface autocorrelations	27
1.16 Standard rotationally invariant fingerprints (RIFs)	29
1.17 Maxima and minima of the local properties	29
1.18 Atom-centered descriptors	29
1.19 Fragment analysis	29
2 PROGRAM OPTIONS	30
2.1 Command-line options	30
2.2 Options defined in the input SDF-file	35
2.2.1 Defining the center for spherical-harmonic fits	35
2.2.2 Defining fragments	36
3 INPUT AND OUTPUT FILES	40
3.1 The VAMP .sdf file as input	41
3.1.1 Multi-structure SD-files	43
3.2 The CeposMopac 6 .sdf file as input	43
3.3 The Vhamil.par file	43



TABLE OF CONTENTS

ParaSurf'12™ Users' Manual

3.4 The EMPIRE™ binary wavefunction file (.vwf)	44
3.5 The ParaSurf™ output file	45
3.5.1 For a spherical-harmonic surface	45
3.5.2 For a marching-cube surface	53
3.5.3 For a job with Shannon entropy	59
3.5.4 For a job with autocorrelation similarity	60
3.6 ParaSurf™ SDF-output	61
3.6.1 Optional blocks in the SDF-output file	64
3.7 The surface (.psf) file	67
3.8 Anonymous SD (.asd) files	68
3.8.1 Optional blocks	69
3.9 Grid calculations with ParaSurf™	70
3.9.1 User-specified Grid	70
3.9.2 Automatic grids	72
3.10 The SIM file format	73
3.11 Output tables	75
3.12 Binned SIM descriptor tables	79
3.13 Autocorrelation fingerprint and similarity tables	80
3.14 Shared files	80
4 TIPS FOR USING PARASURF'12™	81
4.1 Choice of surface	81
4.2 Local properties	81
4.3 QSAR using grids	81
5 SUPPORT	82
5.1 Contact	82
5.2 Error reporting	82
5.3 Cepos InSilico GmbH	82
6 LIST OF TABLES	83
7 LIST OF FIGURES	84
8 REFERENCES	85



PROGRAM HISTORY

Release Date	Version	Platforms
1 st July 2005	ParaSurf'05™ initial release (Revision A1)	32-bit Windows 32-bit Linux Irix
1 st January 2006	ParaSurf'05™ Revision B1 (customer-feedback release)	32-bit Windows 32-bit Linux Irix
1 st July 2006	ParaSurf'06™ Revision A1	32-bit Windows 32-bit Linux 64-bit Linux Irix
1 st July 2007	ParaSurf'07™ Revision A1	
1 st July 2008	ParaSurf'08™ Revision A1	
22 nd August 2008	ParaSurf'08™ Revision A2 (minor bug fix release)	
16 th December 2008	ParaSurf'08™ Revision A3 (minor bug fix release)	
1 st July 2009	ParaSurf'09™ Revision A1	32-bit Windows 64-bit Windows 32-bit Linux 64-bit Linux
1 st September 2009	New Vhamil.par file including PM6 and first-row transition metals in AM1*	
1 st February 2010	ParaSurf'09™ Revision B1 (additional atom-centered descriptors)	
1 st July 2010	ParaSurf'10™ Revision A1	
1 st July 2011	ParaSurf'11™ Revision A1	
1 st September 2013	ParaSurf'12™ Revision A1	



1 INTRODUCTION

ParaSurf™ is a program to generate isodensity or solvent-excluded surfaces from the results of semiempirical molecular orbital calculations, either from VAMP [1] or a public-domain version of MOPAC modified and made available by Cepos InSilico. [2] The surface may be generated by shrink-wrap [3] or marching-cube [4] algorithms and the former may be fit to a spherical harmonic series. [5] The principles of these two techniques are explained below, but for comparison Figure 1 shows default isodensity surfaces calculated by ParaSurf™ for a tetracycline derivative. The surfaces are color-coded according to the electrostatic potential at the surface.




Figure 1 Marching-cube (left) and shrink-wrap (right, fitted to a spherical-harmonic approximation) isodensity surfaces calculated with ParaSurf™ using the default settings

Four local properties, the molecular electrostatic potential (MEP), [6] the local ionization energy (IE_L), [7] the local electron affinity (EA_L), [8] and the local polarizability (α_L) [8] are calculated at the points on the surface. Two further properties, the local hardness (η_L), [8] and the local electronegativity (χ_L) [8] can be derived from IE_L and EA_L .

The local properties can be used to generate a standard set of 81 descriptors [9] appropriate for quantitative structure-property relationships (QSPRs) for determining physical properties.

ParaSurf™ can also generate local enthalpies and free energies of solvation [10] and integrate them over the entire molecular surface to give the enthalpy or free energy of solvation. ParaSurf™ can read so-called *Surface-Integral Model* (SIM) files that allow it to calculate properties such as, for instance, the enthalpy and free energy of hydration and the free energies of solvation in *n*-octanol and chloroform. The surface-integral models are expressed as summations of local solvation energies over the molecular surface. These local solvation energies can be written to the ParaSurf™ surface file.

ParaSurf™ is the first program to emerge from the ParaShift collaboration between researchers at the Universities of Erlangen, Portsmouth, Southampton, Oxford and Aberdeen. It is intended to provide the molecular surfaces for small molecules (*i.e.* non-proteins) for subsequent quantitative structure-activity relationship (QSAR), QSPR, high-throughput virtual screening (HTVS), docking and scoring, pattern-recognition and simulation software that will be developed in the ParaShift project.



1.1 Changes relative to ParaSurf'11™

ParaSurf'12™ has been enhanced relative to its predecessor in order to provide improved flexibility and a more comprehensive range of descriptors and features. The changes are outlined below:

1.1.1 Updated local properties

ParaSurf'11™ and its predecessors used the zero-differential-overlap (ZDO) approximation to calculate electron densities, charges and molecular electrostatics potentials (MEPs) and fields. However, although this approximation is standard in the analysis of neglect of differential diatomic overlap (NDDO) wavefunctions such as those used for ParaSurf™, it does introduce errors. The expressions for the total and orbital electron densities have now been updated in ParaSurf'12™. This gives considerable more accurate results but unfortunately means that the default expressions used in ParaSurf'12™ are no longer compatible with those used in earlier ParaSurf™ versions. We have therefore introduced the command-line option **parasurf11**, which gives backwards compatibility to previous versions of ParaSurf™ by using the old algorithms. Note that ParaSurf'12™ will not give the same results as earlier versions without this option, which must be used for models trained using previous versions of ParaSurf™.

1.1.2 Local polarizability: new definition

The local polarizability used in previous versions of ParaSurf™ (which assumed that atoms are isotropic) has been replaced by a version that takes the anisotropy of the atomic polarizability tensors into account. The older version is also given by the **parasurf11** option.

1.1.3 Electron density: scaling error

A scaling error in the calculated electron density has been corrected. The default isodensity value for the van der Waals-like surface has also been updated. The **parasurf11** option also gives backwards compatibility to older versions in this case.

1.1.4 EMPIRE™

ParaSurf'12™ is compatible with Cepos InSilico's new EMPIRE™ program for performing semiempirical molecular orbital calculations and communicates with EMPIRE™ using the binary **.vWF** file format.

1.1.5 HDF5 files

ParaSurf'13™, which will be released before the end of 2013, will introduce a new HDF5-based binary file communication format to replace ASCII-files such as the SDF file format.



1.2 Isodensity surfaces

Isodensity surfaces [11] are defined as the surfaces around a molecule at which the electron density has a constant value. Usually this value is chosen to approximate the van der Waals' shape of the molecule. ParaSurf™ allows values of the isodensity level down to $0.00001 \text{ e}\cdot\text{\AA}^{-3}$. Lower values than this may result in failures of the surface algorithms for very diffuse surfaces.

1.3 Solvent-excluded surfaces

The solvent-excluded surface is obtained by rolling a spherical solvent molecule of radius r_{solv} over the surface of the molecule as shown in Figure 2. The surface of the solvent molecule defines the molecular surface, so that the yellow volume in Figure 2 becomes part of the molecule.




Figure 2 2D-representation of a solvent-excluded surface

1.4 Solvent-accessible surfaces

Solvent-accessible surfaces are obtained in the same way as solvent-excluded surfaces but the outer surface of the solvent sphere is used to define the molecular surface, as shown in Figure 3.




Figure 3 The solvent-accessible surface is obtained by rolling a spherical "solvent molecule"



1.5 Shrink-wrap surface algorithm

Shrink-wrap surface algorithms [3] are used to determine single-valued molecular surfaces. Single-valued in this case means that for any given radial vector from the center of the molecule the surface is only crossed once (vectors **A** and **B** in Figure 4) and not multiply (vectors **C** and **D** in Figure 4):




Figure 4 2D-representation of a molecular surface with single-valued (A and B) and multiply valued (C and D) radial vectors from the center

Single-valued surfaces are necessary for spherical-harmonic fitting (see [Section 1.4](#)). Thus, spherical-harmonic fitting is only available for shrink-wrap surfaces in ParaSurf™. The shrink-wrap algorithm works by starting outside the molecule (point **a** in Figure 5) and moving inwards along the radial vector until it finds the surface (in our case defined by the predefined level of the electron density, point **b** in Figure 5). Thus, the shrink-wrapped surface may contain areas (marked by dashed lines in Figure 5) for which the surface deviates from the true isodensity surface.

These areas of the surface, however, often have little consequence as they are situated above indentations in the molecule that are poorly accessible to solvents or other molecules. The shrink-wrapped surfaces generated by ParaSurf™ should normally be fitted to a spherical-harmonic series for use in HTVS, similarity, pattern-recognition or high-throughput docking applications. The default molecular center in ParaSurf™ is the center of gravity (CoG). In special cases in which the CoG lies outside the molecule, another center may be chosen.




Figure 5 2D-representation of the shrink-wrap algorithm. The algorithm scans along the vector from point **a** towards the center of the molecule until the electron density reaches the preset value (point **b**). The algorithm results in enclosures (marked yellow) for multi-valued radial vectors



Figure 6 shows a spherical-harmonically fitted shrink-wrap surface for a difficult molecule. The areas shown schematically in Figure 5 are clearly visible.




Figure 6 Spherical-harmonic approximation of a shrink-wrap isodensity surface. Note the areas where the surface does not follow the indentations of the molecule

1.6 Marching-cube algorithm

The marching-cube algorithm [4] implemented in ParaSurf™ does not have the disadvantage of being single-valued like the shrink-wrap surface. It cannot, therefore, be fitted to a spherical harmonic series and is used as a purely numerical surface primarily for QSPR applications or surface-integral models. [10] The algorithm works by testing the electron density at the corners of cubes on a cubic lattice laid out through the molecular volume. The corners are divided into those “inside” the molecule (*i.e.* with a higher electron density than the preset value) and those “outside”. The surface triangulation is then generated for each surface cube and the positions of the surface points corrected to the preset electron density.



1.7 Spherical-harmonic fitting

Complex surfaces can be fitted to spherical harmonic series to give analytical approximations of the surface. [5] The surfaces are fit to a series of distances $r_{\alpha,\beta}$ from the center along the radial vector defined by the angles α and β as:

$$r_{\alpha,\beta} = \sum_{l=0}^N \sum_{m=-l}^l c_l^m Y_l^m \quad (1)$$

Where the distances $r_{\alpha,\beta}$ are linear combinations of spherical harmonics Y_l^m defined as:

$$Y_l^m(\alpha, \beta) = \sqrt{\frac{(2l+1)(l-m)!}{4\pi(l+m)!}} P_l^m(\cos \alpha) e^{im\beta} \quad (2)$$

where $P_l^m(\cos \alpha)$ are associated Legendre functions and l and m are integers such that $-l \leq m \leq l$. In the above form, spherical harmonics are complex functions. Duncan and Olson [12] have used the real functions

$$Y_l^m(\alpha, \beta) = N_{lm} P_l^m(\cos \alpha) \cos |m| \beta \quad (3)$$

where N_{lm} are normalization factors, to describe molecular surfaces using spherical harmonics.

ParaSurf™ not only fits the surface itself (i.e. the radial distances) to spherical harmonic expansions, but also the four local properties (see [Section 1.8](#)). In this way, a completely analytical description of the shape of the molecule and its intermolecular binding properties is obtained. [13] This description can be truncated at different orders l depending on the application and the precision needed. Thus, a simple description of the molecular properties (shape, MEP, IE_L, EA_L and α_L) to order 2 consists of only five sets of nine coefficients each, or 45 coefficients. These coefficients can be rotated, overlaps calculated etc. [5] to give fast scanning of large numbers of compounds.

Note that, because of the approximate nature of the spherical-harmonic fits, the default isodensity level for the shrink-wrapped surface ($0.0005 \text{ e}\cdot\text{\AA}^{-3}$) is lower than that ($0.007 \text{ e}\cdot\text{\AA}^{-3}$) appropriate for an approximately van der Waals' surface using the marching-cube algorithm. The lower value avoids the surface coming too close to atoms. Note also that the fits are incremental, which means that the order chosen for a given application can be obtained by ignoring coefficients of higher order in the spherical-harmonic series.

In some cases, the default resolution of the molecular surface does not allow fitting the spherical-harmonic expansion to very high orders without introducing noise ("ripples") on the fitted surface. In this case, the calculated RMSD becomes larger at higher orders of the spherical-harmonic expansion. ParaSurf'12™ recognizes this condition and truncates the fitting procedure at the optimum value. This can be recognized in the output because the RMSD for later cycles remains constant and the coefficients of the higher order spherical harmonics are all zero. This guarantees the optimum fit in each case and is important for applications that use either the spherical-harmonic coefficients themselves or the hybridization coefficients.



The choice of center for fitting to a spherical-harmonic expansion is critical. ParaSurf'12™ therefore goes through a multi-step procedure in order to find a suitable center. This procedure is retained for all molecules for which the ParaSurf'08™ found a suitable center. However, if the algorithms implemented in ParaSurf'08™ fail to find a suitable center, the additional technique implemented in ParaSurf'12™ will probably work.

The problem with many molecules is that, for instance, the center of mass does not lie within the molecular volume. This can easily be the case for, for instance, U- or L-shaped molecules. The procedure implemented in ParaSurf'12™ works as follows:

1. The program first calculates the center of mass and tests whether it lies within the volume of the molecule. If it does, it is used as the molecular center. If not, the program moves on to the next step.
2. ParaSurf™ calculates the principal moments of inertia of the molecule and derives a center from them by assuming that the molecule is U- or V-shaped. The procedure tries to place the center at the base center of the molecule. This procedure was implemented in ParaSurf'08™ as a fallback if the center of mass proved unsuitable. If it also fails to find a suitable center, ParaSurf'12™ moves on to a third option, which finds a center for all but the most difficult molecules.
3. The new procedure first searches for the largest plane in the molecule (*i.e.* the one that contains the most atoms). This search has some leeway, so that the atoms must not all lie exactly in the plane. As a second step, the second largest plane is sought. The molecular center is then placed in the hinge area between the two planes, as illustrated in Figure 7:




Figure 7 Schematic representation of the planes and hinge area used to determine the center for spherical-harmonic expansions



1.8 Local properties

The local properties calculated by ParaSurf™ are those related to intermolecular interactions. Local properties, sometimes inaccurately called fields in QSAR work, are properties that vary in space around the molecule and therefore have a distribution of values at the molecular surface. The best known and most important local property in this context is the molecular electrostatic potential, which governs Coulomb interactions, but the MEP only describes a part of the intermolecular interaction energy, so that further local properties are needed.

1.8.1 Molecular electrostatic potential

The MEP is defined in ParaSurf™ as the energy of interaction of a single positive electronic charge at the position \mathbf{r} with the molecule. Within quantum mechanical (semiempirical or *ab initio* molecular orbital (MO) theory, density functional theory (DFT)) the MEP ($V(\mathbf{r})$) is described [6] as:

$$MEP(\mathbf{r}) = \sum_{i=1}^n \frac{Z_i}{|\mathbf{R}_i - \mathbf{r}|} - \int \frac{\rho(\mathbf{r}') d\mathbf{r}'}{|\mathbf{r}' - \mathbf{r}|} \quad (4)$$

where n is the number of atoms in the molecule, Z_i is the nuclear charge of atom i located at \mathbf{R}_i , and $\rho(\mathbf{r})$ is the electron-density function of the molecule. This expression, however, involves integrating the electron density, a time-consuming calculation. ParaSurf™ therefore uses two different approximate models for calculating the MEP.

1.8.1.1 The natural atomic orbital/PC (NAO-PC) model

The NAO-PC model [14] uses a total of nine point charges, one positive charge at the nucleus and eight negative ones distributed around it, to describe the electrostatics of a non-hydrogen atom with a valence-only *s*- and *p*-basis set for the semiempirical Hamiltonians MNDO, [15] AM1 [16] and PM3 [17]. The negative charges are located at the charge centers of each lobe of the natural atomic orbitals, which are obtained by diagonalizing the one-atom blocks of the density matrix. [18] The NAO-PC charges are calculated by VAMP and output in the `.sdf` file for use in ParaSurf™. The NAO-PC model is therefore only available when using ParaSurf™ with VAMP `.sdf` input. NAO-PC charges are also not available for semiempirical Hamiltonians such as MNDO/d [19] or AM1* [18] that use *d*-orbitals in the basis set.

1.8.1.2 The multipole model

The integrals needed to evaluate Eq. (4) in MNDO-type methods use a multipole approximation [20] that extends to quadrupoles. We can therefore also use this approximation to calculate atom-centered monopoles, dipoles and quadrupoles for each atom in the molecule. [21] This multipole model is applicable to all methods, including those with *d*-orbitals, and can be used with MOPAC output files as input to ParaSurf™.



1.8.2 Local ionization energy, electron affinity, hardness and electronegativity

The local ionization energy $IE_L(\mathbf{r})$ is defined [7] as a density-weighted Koopmans' ionization potential at a point \mathbf{r} near the molecule:

$$IE_L(\mathbf{r}) = \frac{-\sum_{i=1}^{HOMO} \rho_i(\mathbf{r}) \varepsilon_i}{\sum_{i=1}^{HOMO} \rho_i(\mathbf{r})} \quad (5)$$

where $HOMO$ is the number of the highest occupied MO, $\rho_i(\mathbf{r})$ is the electron density at point \mathbf{r} due to MO i and ε_i is its Eigenvalue. The local ionization energy describes the tendency of the molecule to interact with electron acceptors (Lewis acids) in a given region in space. [7-8]

The definition of the local electron affinity is a simple extension of [Eq. \(5\)](#) to the virtual MOs: [8]

$$EA_L(\mathbf{r}) = \frac{-\sum_{i=LUMO}^{norbs} \rho_i(\mathbf{r}) \varepsilon_i}{\sum_{i=LUMO}^{norbs} \rho_i(\mathbf{r})} \quad (6)$$

The local electron affinity is the equivalent of the local ionization energy for interactions with electron donors (Lewis bases). [8] An intensity-filtering technique [20a] was introduced in ParaSurf'10™ to allow the local electron affinity to be calculated for Hamiltonians such as AM1* and MNDO/d that use polarization d -functions.

Equation (6) requires that the occupied and virtual orbitals be approximately equivalent to each other. This is not the case for semiempirical Hamiltonians (such as AM1*) that include d -orbitals as polarization functions or for extensive basis sets in Hartree-Fock *ab initio* or in Density-Functional theory (DFT) calculations. A new technique has therefore been defined [11] to exclude pure polarization functions from the summation in [Eq. \(6\)](#). This technique is now default in ParaSurf'12™ and gives reliable results. For continuity, a new command-line option (EAL09) has been introduced to request that the calculation of the local electron affinity be performed exactly as in ParaSurf'09™ and earlier versions.

Two further, less fundamental local properties have been defined. [8] These are the local hardness, η_L :

$$\eta_L = \frac{(IE_L - EA_L)}{2} \quad (7)$$

and the local electronegativity, χ_L :

$$\chi_L = \frac{(IE_L + EA_L)}{2} \quad (8)$$



1.8.3 Local polarizability

Within the NDDO, the molecular electronic polarizability is easily accessible using the parameterized version [22] of the variational technique introduced by Rivail, [23] which can also be partitioned into an additive polarizability scheme. [20b] Versions of ParaSurf™ up to ParaSurf'11™ used an isotropic definition of the local polarizability, α_L , at a point near the molecule:

$$\alpha_L(\mathbf{r}) = \frac{\sum_{j=1}^{norbs} \rho_j^1(\mathbf{r}) q_j \bar{\alpha}_j}{\sum_{j=1}^{norbs} \rho_j^1(\mathbf{r}) q_j} \quad (9)$$

where q_j is the Coulson occupation and $\bar{\alpha}_j$ the isotropic polarizability attributed to atomic orbital j . The density ρ_j^1 is defined as the electron density at the point in question due to an exactly singly occupied atomic orbital j . The sum is now over atomic orbitals, rather than MOs as for the other local properties. Thus, the local polarizability is a simple occupation-weighted sum of the orbital polarizabilities in which the contribution of each AO is determined by the density of the individual AO at the point being considered.

ParaSurf'12™ makes use of the fact that the atomic polarizability tensors produced by the procedure described in reference [20b] are anisotropic. It uses this atomic anisotropy to calculate a more highly resolved local polarizability that is now standard in ParaSurf'12™. The keyword **parasurf11** ensures backwards compatibility with the isotropic local polarizability used in earlier versions.

1.8.4 Field normal to the surface

The electrostatic field (the first derivative of the potential) normal to the molecular surface is closely related to the electrostatic solvation energy in implicit solvation models. [24] This field also has the advantage that it is largely independent of the total molecular charge, so that charged molecules can be compared with neutral ones. If the molecular electrostatic potential is used for this purpose, the charge of ions leads a shift in the potential descriptors, so that molecules and ions with different charges cannot be compared directly. The direction of the normal field (inwards or outwards) also defines, for instance hydrogen-bond donors and acceptors specifically.



1.9 Descriptors

A set of 81 molecular descriptors derived from the MEP, local ionization energy, IE_L , electron affinity, EA_L , electronegativity, χ_L , hardness, η_L , and polarizability, α_L has been defined for QSPR-studies. [9] These and several related descriptors calculated and output by ParaSurf™ are defined in the following table (Table 1).

Table 1 The descriptors calculated by ParaSurf™

Descriptor	Description	Formula/ Reference	Symbol in CSV file
μ	Dipole moment		dipole
μ_D	Dipolar density	[20b]	dipden
α	Molecular electronic polarizability	[25]	polarizability
MW	Molecular weight		Mwt
G	Globularity	[26]	globularity
A	Molecular surface area		totalarea
VOL	Molecular volume		volume
V_{\max}	Maximum (most positive) MEP	[27]	MEPmax
V_{\min}	Minimum (most negative) MEP	[27]	MEPmin
\bar{V}_+	Mean of the positive MEP values	[27]	meanMEP+
\bar{V}_-	Mean of the negative MEP values	[27]	meanMEP-
\bar{V}	Mean of all MEP values	[27]	meanMEP
ΔV	MEP-range	[27]	MEP-range
σ_+^2	Total variance in the positive MEP values	[27]	MEPvar+
σ_-^2	Total variance in the negative MEP values	[27]	MEPvar-
σ_{tot}^2	Total variance in the MEP	[27]	MEPvartot
v	MEP balance parameter	[27]	MEPbalance
$\sigma_{tot}^2 v$	Product of the total variance in the MEP and the balance parameter	[27]	var*balance
γ_1^V	Skewness of the MEP-distribution	$\gamma_1^{\alpha_L} = \frac{\sum_{i=1}^N (\alpha_L^i - \bar{\alpha}_L)^3}{(N-1)\sigma^3}$	MEPskew



Descriptor	Description	Formula/ Reference	Symbol in CSV file
γ_2^V	Kurtosis of the MEP-distribution	$\gamma_2^V = \frac{\sum_{i=1}^N (V_i - \bar{V})^4}{(N-1)\sigma^4} - 3$	MEPkurt
\int_V	Integrated MEP over the surface	$\int_V = \sum_{i=1}^N V_i a_i$	MEPint
IE_L^{\max}	Maximum value of the local ionization energy		IELmax
IE_L^{\min}	Minimum value of the local ionization energy		IELmin
\overline{IE}_L	Mean value of the local ionization energy	$\overline{IE}_L = \frac{1}{N} \sum_{i=1}^N IE_L^i$	IELbar
ΔIE_L	Range of the local ionization energy	$\Delta IE_L = IE_L^{\max} - IE_L^{\min}$	IELrange
σ_{IE}^2	Variance in the local ionization energy	$\sigma_{IE}^2 = \frac{1}{N} \sum_{i=1}^N [IE_L^i - \overline{IE}_L]^2$	IELvar
$\gamma_1^{IE_L}$	Skewness of the local ionization energy distribution	$\gamma_1^{IE_L} = \frac{\sum_{i=1}^N (IE_L^i - \overline{IE}_L)^3}{(N-1)\sigma^3}$	IELSKEW
$\gamma_2^{IE_L}$	Kurtosis of the local ionization energy distribution	$\gamma_2^{IE_L} = \frac{\sum_{i=1}^N (IE_L^i - \overline{IE}_L)^4}{(N-1)\sigma^4} - 3$	IELkurt
\int_{IE_L}	Integrated local ionization energy over the surface	$\int_{IE_L} = \sum_{i=1}^N IE_L^i a_i$	IELint
EA_L^{\max}	Maximum of the local electron affinity		EALmax
EA_L^{\min}	Minimum of the local electron affinity		EALmin
\overline{EA}_{L+}	Mean of the positive values of the local electron affinity	$\overline{EA}_{L+} = \frac{1}{N^+} \sum_{i=1}^{N^+} EA_{L+}^i$	EALbar+
\overline{EA}_{L-}	Mean of the negative values of the local electron affinity	$\overline{EA}_{L-} = \frac{1}{N^-} \sum_{i=1}^{N^-} EA_{L-}^i$	EALbar-
\overline{EA}_L	Mean value of the local electron affinity	$\overline{EA}_L = \frac{1}{N} \sum_{i=1}^N EA_L^i$	EALbar
ΔEA_L	Range of the local electron affinity	$\Delta EA_L = EA_L^{\max} - EA_L^{\min}$	EALrange
σ_{EA+}^2	Variance in the local electron affinity for all positive values	$\sigma_{EA+}^2 = \frac{1}{m} \sum_{i=1}^m [EA_i^+ - \overline{EA}^+]^2$	EALvar+



Descriptor	Description	Formula/ Reference	Symbol in CSV file
$\sigma_{EA^-}^2$	Variance in the local electron affinity for all negative values	$\sigma_{EA^-}^2 = \frac{1}{n} \sum_{i=1}^n \left[EA_i^- - \bar{EA}^- \right]^2$	EALvar-
$\sigma_{EA_{tot}}^2$	Sum of the positive and negative variances in the local electron affinity	$\sigma_{EA_{tot}}^2 = \sigma_{EA^+}^2 + \sigma_{EA^-}^2$	EALvartot
v_{EA}	Local electron affinity balance parameter	$v_{EA} = \frac{\sigma_{EA^+}^2 \cdot \sigma_{EA^-}^2}{\left[\sigma_{EA}^2 \right]^2}$	EALbalance
δA_{EA}^+	Fraction of the surface area with positive local electron affinity	$\delta A_{EA}^+ = \frac{A_{EA}^+}{A},$ A = total surface area	EALfraction+
A_{EA}^+	Surface area with positive local electron affinity		EALarea+
$\gamma_1^{EA_L}$	Skewness of the local electron affinity distribution	$\gamma_1^{EA_L} = \frac{\sum_{i=1}^N \left(EA_L^i - \bar{EA}_L \right)^3}{(N-1)\sigma^3}$	EALskew
$\gamma_2^{EA_L}$	Kurtosis of the local electron affinity distribution	$\gamma_2^{EA_L} = \frac{\sum_{i=1}^N \left(EA_L^i - \bar{EA}_L \right)^4}{(N-1)\sigma^4} - 3$	EALKurt
\int_{EA_L}	Integrated local electron affinity over the surface	$\int_{IE_L} = \sum_{i=1}^N EA_L^i a_i$	EALint
α_L^{\max}	Maximum value of the local polarizability		POLmax
α_L^{\min}	Minimum value of the local polarizability		POLmin
$\bar{\alpha}_L$	Mean value of the local polarizability	$\bar{\alpha}_L = \frac{1}{N} \sum_{i=1}^N \alpha_L^i$	POLbar
$\Delta \alpha_L$	Range of the local polarizability	$\Delta \alpha_L = \alpha_L^{\max} - \alpha_L^{\min}$	POLrange
σ_{α}^2	Variance in the local polarizability	$\sigma_{\alpha}^2 = \frac{1}{N} \sum_{i=1}^N \left[\alpha_L^i - \bar{\alpha}_L \right]^2$	POLvar
$\gamma_1^{\alpha_L}$	Skewness of the local polarizability distribution	$\gamma_1^{\alpha_L} = \frac{\sum_{i=1}^N \left(\alpha_L^i - \bar{\alpha}_L \right)^3}{(N-1)\sigma^3}$	POLskew
$\gamma_2^{\alpha_L}$	Kurtosis of the local polarizability distribution	$\gamma_2^{\alpha_L} = \frac{\sum_{i=1}^N \left(\alpha_L^i - \bar{\alpha}_L \right)^4}{(N-1)\sigma^4} - 3$	POLkurt
\int_{α_L}	Integrated local polarizability over the surface	$\int_{\alpha_L} = \sum_{i=1}^N \alpha_L^i a_i$	POLint



Descriptor	Description	Formula/ Reference	Symbol in CSV file
χ_L^{\max}	Maximum value of the local electronegativity		ENEGmax
χ_L^{\min}	Minimum value of the local electronegativity		ENEGmin
$\bar{\chi}_L$	Mean value of the local electronegativity	$\bar{\chi}_L = \frac{1}{N} \sum_{i=1}^N \chi_L^i$	ENEGbar
$\Delta\chi_L$	Range of the local electron electronegativity	$\Delta\chi_L = \chi_L^{\max} - \chi_L^{\min}$	ENEGrange
σ_χ^2	Variance in the local electronegativity	$\sigma_\chi^2 = \frac{1}{N} \sum_{i=1}^N \left[\chi_L^i - \bar{\chi}_L \right]^2$	ENEGvar
$\gamma_1^{\chi_L}$	Skewness of the local electronegativity distribution	$\gamma_1^{\chi_L} = \frac{\sum_{i=1}^N (\chi_L^i - \bar{\chi}_L)^3}{(N-1)\sigma^3}$	ENEGskew
$\gamma_2^{\chi_L}$	Kurtosis of the local electronegativity distribution	$\gamma_2^{\chi_L} = \frac{\sum_{i=1}^N (\chi_L^i - \bar{\chi}_L)^4}{(N-1)\sigma^4} - 3$	ENEGkurt
\int_{χ_L}	Integrated local electronegativity over the surface	$\int_{\chi_L} = \sum_{i=1}^N \chi_L^i a_i$	ENEGint
η_L^{\max}	Maximum value of the local hardness		HARDmax
η_L^{\min}	Minimum value of the local hardness		HARDmin
$\bar{\eta}_L$	Mean value of the local hardness	$\bar{\eta}_L = \frac{1}{N} \sum_{i=1}^N \eta_L^i$	HARDbar
$\Delta\eta_L$	Range of the local electron hardness	$\Delta\eta_L = \eta_L^{\max} - \eta_L^{\min}$	HARDrange
σ_η^2	Variance in the local hardness	$\sigma_\eta^2 = \frac{1}{N} \sum_{i=1}^N \left[\eta_L^i - \bar{\eta}_L \right]^2$	HARDvar
$\gamma_1^{\eta_L}$	Skewness of the local hardness distribution	$\gamma_1^{\eta_L} = \frac{\sum_{i=1}^N (\eta_L^i - \bar{\eta}_L)^3}{(N-1)\sigma^3}$	HARDskew
$\gamma_2^{\eta_L}$	Kurtosis of the local hardness distribution	$\gamma_2^{\eta_L} = \frac{\sum_{i=1}^N (\eta_L^i - \bar{\eta}_L)^4}{(N-1)\sigma^4} - 3$	HARDkurt
\int_{η_L}	Integrated local hardness over the surface	$\int_{\eta_L} = \sum_{i=1}^N \eta_L^i a_i$	HARDint
F_N^{\max}	Maximum value of the electrostatic field normal to the surface		FNmax



Descriptor	Description	Formula/ Reference	Symbol in CSV file
F_N^{\min}	Minimum value of the field normal to the surface		FNmin
\bar{F}_N	Mean value of the field normal to the surface	$\bar{F}_N = \frac{1}{N} \sum_{i=1}^N \chi_L^i$	FNmean
σ_F^2	Variance in field normal to the surface	$\sigma_F^2 = \frac{1}{N} \sum_{i=1}^N \left[F_N^i - \bar{F}_N \right]^2$	FNvartot
σ_{F+}^2	Variance in the field normal to the surface for all positive values	$\sigma_{F+}^2 = \frac{1}{m} \sum_{i=1}^m \left[F_N^{i+} - \bar{F}_N^+ \right]^2$	FNvar+
σ_{F-}^2	Variance in the field normal to the surface for all negative values	$\sigma_{F-}^2 = \frac{1}{n} \sum_{i=1}^n \left[F_N^{i-} - \bar{F}_N^- \right]^2$	FNvar-
v_F	Normal field balance parameter	$v_F = \frac{\sigma_{F+}^2 \cdot \sigma_{F-}^2}{\left[\sigma_F^2 \right]^2}$	FNbal
$\gamma_1^{F_N}$	Skewness of the field normal to the surface	$\gamma_1^{F_N} = \frac{\sum_{i=1}^N \left(F_N^i - \bar{F}_N \right)^3}{(N-1)\sigma^3}$	FNskew
$\gamma_2^{F_N}$	Kurtosis of the field normal to the surface	$\gamma_2^{F_N} = \frac{\sum_{i=1}^N \left(F_N^i - \bar{F}_N \right)^4}{(N-1)\sigma^4} - 3$	FNkurt
\int_{F_N}	Integrated field normal to the surface over the surface	$\int_{F_N} = \sum_{i=1}^N F_N^i a_i$	FNint
$\int_{F_N}^+$	Integrated field normal to the surface over the surface for all positive values	$\int_{F_N}^+ = \sum_{i=1}^N F_N^i a_i \text{ if } F_N^i \geq 0$	FN+
$\int_{F_N}^-$	Integrated field normal to the surface over the surface for all negative values	$\int_{F_N}^- = \sum_{i=1}^N F_N^i a_i \text{ if } F_N^i < 0$	FN-
$\int_{ F_N }$	Integrated absolute field normal to the surface over the surface	$\int_{ F_N } = \sum_{i=1}^N F_N^i a_i$	FNabs
Additionally if the Shannon Entropy is calculated			
H_{in}^{\max}	Maximum value of the internal Shannon Entropy		SHANImax
H_{in}^{\min}	Minimum value of the internal Shannon Entropy		SHANImin
\bar{H}_{in}	Mean value of the internal Shannon Entropy	$\bar{H}_{in} = \frac{1}{N} \sum_{i=1}^N H_{in}^i$	SHANIBar
$\sigma_{H_{in}}^2$	Variance in the internal Shannon Entropy	$\sigma_{H_{in}}^2 = \frac{1}{N} \sum_{i=1}^N \left[H_{in}^i - \bar{H}_{in} \right]^2$	SHANIVar



Descriptor	Description	Formula/ Reference	Symbol in CSV file
$\int_{H_{in}}$	Integrated internal Shannon Entropy over the surface	$\int_{H_{in}} = \sum_{i=1}^N H_{in}^i a_i$	SHANItot
And if the external Shannon Entropy is available			
H_{ex}^{\max}	Maximum value of the external Shannon Entropy		SHANEmax
H_{ex}^{\min}	Minimum value of the external Shannon Entropy		SHANEmin
$\overline{H_{ex}}$	Mean value of the external Shannon Entropy	$\overline{H_{ex}} = \frac{1}{N} \sum_{i=1}^N H_{ex}^i$	SHANEbar
$\sigma_{H_{ex}}^2$	Variance in the external Shannon Entropy	$\sigma_{H_{ex}}^2 = \frac{1}{N} \sum_{i=1}^N [H_{ex}^i - \overline{H_{ex}}]^2$	SHANEvar
$\int_{H_{ex}}$	Integrated internal Shannon Entropy over the surface	$\int_{H_{ex}} = \sum_{i=1}^N H_{ex}^i a_i$	SHANEtot

1.10 Surface-integral models (polynomial version)

The polynomial surface-integral models that can be calculated by ParaSurf™ are defined [10] using the expression

$$P = \sum_{i=1}^{ntri} f(V^i, IE_L^i, EA_L^i, \alpha_L^i, \eta_L^i) \cdot A^i \quad (10)$$

where P is the target property, usually a free energy, f is a polynomial function of the electrostatic potential V , the local ionization energy, IE_L , the local electron affinity, EA_L , the local polarizability, α_L and the local hardness, η_L . A^i is the area of the surface triangle i .

The molecular property P is printed to the output file and to the `<filename>.p.sdf` ParaSurf™ output SD-file. The individual values of the function f are added to the list of local properties written for each surface point to the `.psf` file if the surface details are output.

The surface-integral models themselves are not implemented directly in ParaSurf™, but are read in general form from the SIM file, whose format is given in [Section 3.10](#). Thus, the users' own surface-integral models can be added to ParaSurf™. Data for generating surface-integral models can be derived simply from the `.psf` surface output for a normal ParaSurf™ run. Note that the program options given in the SIM file must be the same for all the models included in the file and that they override conflicting command-line options.



1.11 Binned surface-integral models

A more recent type of SIM model, binned SIM models, [13] is now implemented in ParaSurf'12™ for the negative logarithm of the water-octanol partition constant, logPow. These models divide the surface into bins according to the values of the local properties and use the total surface area assigned to each bin as descriptors for multiple linear regression models. These models have been implemented for marching cube surfaces using either the isodensity or solvent-excluded surfaces and for the AM1, AM1*, MNDO, MNDO/d, PM3 and PM6 Hamiltonians. In contrast to polynomial SIM models, they are encoded in the program and are output under the heading **ParaSurf ADMET Profiler**. These logPow models are available for the MNDO, AM1, PM3, MNDO/d, AM1* and PM6 Hamiltonians. The models use “conformationally averaged” structures within a standard calculational protocol in which the initial 3D structure is produced by CORINA [28] as the starting geometry for the semiempirical geometry optimization and uses only this one conformation to predict logPow for the compound. These models were trained with all verified values contained in the LogKOW dataset [29] and are those denoted “single conformation” trained with the “full” dataset in the original literature. [30]

Table 2 The 28 local properties and products thereof used to construct binned area descriptors.

	IE_L	EA_L	α_L^a	F_N	η_L	χ_L
MEP	$MEP \times IE_L$	$MEP \times EA_L$	$MEP \times \alpha_L$	$MEP \times F_N$	$MEP \times \eta_L$	$MEP \times \chi_L$
IE_L		$IE_L \times EA_L$	$IE_L \times \alpha_L$	$IE_L \times F_N$	$IE_L \times \eta_L$	$IE_L \times \chi_L$
EA_L			$EA_L \times \alpha_L$	$EA_L \times F_N$	$EA_L \times \eta_L$	$EA_L \times \chi_L$
α_L^a				$\alpha_L \times F_N$	$\alpha_L \times \eta_L$	$\alpha_L \times \chi_L$
F_N					$F_N \times \eta_L$	$F_N \times \chi_L$
η_L						$\eta_L \times \chi_L$

^a Not used for MNDO/d, AM1* or PM6

Local hydrophobicities and logPow models are available for the following combinations of Hamiltonians, surfaces and contours. The three letter model code is used to write the local hydrophobicity to the output .vmp and .psf files or to specify that the descriptors for the model are written out.

Table 3 Local hydrophobicity models and their model codes (all models use the single CORINA-derived conformations and are trained with the “full” dataset. [30])

Hamiltonian	Model code	
	ParaSurf'11™	ParaSurf'12™
AM1	LP1	OW1
AM1*	LP2	OW2
PM3	LP3	OW3
MNDO	LP4	OW4
MNDO/d	LP5	OW5
PM6	LP6	OW6



1.12 Spherical harmonic “hybrids”

Once the molecular shape or a local property have been fitted to a spherical-harmonic expansion, [16] the shape or property can be described succinctly as a series of spherical-harmonic “hybridization” coefficients analogous to the concept of hybrid atomic orbitals. Thus, for each value of l in [Eq. \(1\)](#) the “hybridization” coefficient H_l is given by:

$$H_l = \sum_{m=-l}^l \left(c_l^m \right)^2 \quad (11)$$

The hybridization coefficients H_l can be used as additional descriptors for fast QSPR screening.



1.13 Descriptors and moments based on polynomial surface-integral models

ParaSurf™ uses local properties defined in a surface-integral model (SIM, see [Section 1.10](#)) to calculate descriptors analogous to those listed in [Table 1](#). Additionally, **dipolarmoments** of the local property are calculated. These are gauge-independent moments calculated by first shifting values of the local property so that their sum is zero and then calculating moments according to

$$\mu = \sum_{i=1}^{ntri} P_i \mathbf{r}_i \quad (12)$$

where μ is the dipolar moment, P_i the value of the local property i situated at position \mathbf{r}_i .

The output for these properties derived from a SIM for logPow is shown below:

Descriptors calculated for logP:

Dipolar moment	x:	-549.2	y:	-247.9	z:	-937.0
Sum:	1114.					
Most positive value	:	1.407				
Most negative value	:	0.8325E-01				
Range	:	1.324				
Mean	:	0.1874				
Mean positive	:	0.1874				
Mean negative:		0.000				
Total variance:		0.2376E-01				
Positive variance:		0.2376E-01				
Negative variance	:	0.000				
Balance parameter	:	0.000				
Balance*variance	:	0.000				

The values of these descriptors are often useful for deriving models directly related to the property modelled by the SIM. Note that no units are given in the output because they depend on the property modelled by the SIM.



1.14 Shannon entropy

The information content at the surface of the molecule can be defined based on the distribution of the four local properties over the surface using an approach analogous to that introduced by Shannon. [31]

Shannon defined the Shannon entropy, H , which corresponds to the amount of information (in bits) as

$$H = -\sum_{i=1}^n p_i \log_2(p_i) \quad (13)$$

where n is the number of possible characters and p_i is the probability that character i will occur. Note that, importantly, this definition of the amount of information is local (*i.e.* it only depends on the value of the probability of character i).

For a continuous property, X , Eq. (13) becomes

$$H = -\int_{-\infty}^{\infty} p(X) \log_2 p(X) dX \quad (14)$$

If we now assume that the Shannon entropy at a point in space near a molecule is defined by the values of the four continuous local properties described above, we obtain

$$H = -\iiint p(V, I, E, \alpha) \log_2 p(V, I, E, \alpha) dV dI dE d\alpha \quad (15)$$

where $p(V, I, E, \alpha)$ is the probability of finding the values V, I, E and α . However, we can simplify this expression because the four properties are essentially independent of each other, [8-9] so that we can write

$$\begin{aligned} H = & -\int p(V) \log_2 p(V) dV - \int p(I) \log_2 p(I) dI \\ & - \int p(E) \log_2 p(E) dE - \int p(\alpha) \log_2 p(\alpha) d\alpha \end{aligned} \quad (16)$$

Transferring this definition to a molecule for which a triangulated surface of k triangles, where triangle i has area A_i and average values of the four local properties V_i, I_i, E_i and α_i we obtain

$$H = -\sum_{i=1}^k [p(V_i) \log_2 p(V_i) + p(I_i) \log_2 p(I_i) + p(E_i) \log_2 p(E_i) + p(\alpha_i) \log_2 p(\alpha_i)] \cdot A_i \quad (17)$$

where $p(X_i)$ is the probability that the value X_i of the property X , where X may be V, I, E or α , will occur.

ParaSurf™ offers two alternatives as sources for the probabilities $p(X_i)$. The first, known as the “external” Shannon entropy, is to use probabilities taken from an external dataset and defined in a separate statistics file. The default “external” statistics file is called **bins.txt** and is read from the



ParaSurf™ root directory. The statistics defined in **bins.txt** were derived from AM1 calculations of all the bound ligands defined in the PDBbind database [32] in their correct protonation states and at geometries obtained by optimizing with AM1 starting from the bound conformation. [27]

Alternatively, the user can define a custom “external” statistics file using the ParaSurf™ module **binner** (available free of charge for ParaSurf™ users). The “external” Shannon entropy is useful for relating a series of molecules to each other, but is sensitive, for instance, to the total charge of the molecule.

The “internal” Shannon entropy is calculated using probabilities determined from the surface properties of the molecule itself, and therefore corresponds more closely to Shannon’s classical definition than the “external” Shannon entropy and the probabilities used are individual for each molecule. The “internal” Shannon entropy can be considered to represent the information content of the molecule. The properties of the two types of Shannon entropy will be described in a forthcoming paper.



1.15 Surface autocorrelations

Gasteiger *et al.* [33] introduced the concept of surface autocorrelations as powerful descriptions of molecular binding properties for quantitative structure-activity relationships (QSARs). In ParaSurf™, autocorrelations $A(R)$ are now defined as:

$$A(R) = \sum_{i=1}^{\text{npoints}-1} \sum_{j=i+1}^{\text{npoints}} \omega_{ij} \delta(R, r_{ij}) \quad (18)$$

where r_{ij} is the distance between surface points i and j and ω_{ij} is a function of one or more local properties at the points i and j and δ_{ij} is 1.0 if r_{ij} is inside the bin centered on R and zero otherwise. Note that this is a different definition of the autocorrelation function to that used in earlier versions of ParaSurf™. Also, because the new algorithm for calculating the autocorrelations is very fast, all surface points are used, rather than sampling 10% as earlier.

Seven different autocorrelation functions are calculated by ParaSurf™. These are:

Shape autocorrelation	$\omega_{ij} = 1.0$
MEP autocorrelation	$\omega_{ij} = V_i \times V_j$
Plus-plus MEP autocorrelation	$\omega_{ij} = V_i \times V_j$ $\omega_{ij} = 0.0$ ($V_i > 0$ and $V_j > 0$) ($V_i < 0$ or $V_j < 0$)
Minus-minus MEP autocorrelation	$\omega_{ij} = V_i \times V_j$ ($V_i < 0$ and $V_j < 0$)
Plus-minus MEP autocorrelation	$\omega_{ij} = -V_i \times V_j$ $\omega_{ij} = 0.0$ ($V_i \times V_j < 0$) ($V_i \times V_j > 0$)
Local ionization energy autocorrelation	$\omega_{ij} = IE_L^i \times IE_L^j$
Local electron affinity autocorrelation	$\omega_{ij} = EA_L^i \times EA_L^j$
Local polarizability autocorrelation	$\omega_{ij} = \alpha_L^i \times \alpha_L^j$

Generally, the shape autocorrelation and that based on the local polarizability correlate strongly with each other. The MEP correlation is the sum of its three components (plus-plus, plus-minus and minus-minus). However the three components enable us to distinguish between ++ and – pairs of surface points, which both give a positive contribution to the autocorrelation function.



ParaSurf™ calculates autocorrelations as vectors of $A(R)$ values 64 elements long starting at an R -value of 0.0 Å and increasing in bins of width 0.2 Å up to a maximum value of 12.8 Å). Figure 8 shows the eight autocorrelation functions for trimethoprim calculated with AM1.




Figure 8 The four autocorrelation functions calculated using the AM1 Hamiltonian for trimethoprim

The command-line argument `autocorr=<filename>` requests that similarities in the autocorrelation functions with the molecule described in `<filename>`, where `<filename>` must be a `ParaSurf.sdf` output file, are calculated and written out. The floating-point Tanimoto similarities S are defined as:

$$S = \frac{\sum_{i=1}^{nbins} A_A^i A_B^i}{\sum_{i=1}^{nbins} (A_A^i)^2 + \sum_{i=1}^{nbins} (A_B^i)^2 - \sum_{i=1}^{nbins} A_A^i A_B^i} \quad (19)$$

where A_A^i is the value of the autocorrelation function for molecule A and bin i etc. Note that although the normal range for a Tanimoto coefficient is from zero to one, marginally negative values may occur for the local ionization energy or electron affinity.

These similarities are calculated for the entire range of each of the eight autocorrelation functions. These individual similarities can be written to a table file (see [Section 3.13](#)) and are printed in the output file (see [Section 3.5.4](#)).



1.16 Standard rotationally invariant fingerprints (RIFs)

Mavridis *et al.* [34] introduced standard rotationally invariant fingerprints (RIFs) based on the spherical-harmonic hybridization coefficients defined above. These fingerprints provide a detailed description of the molecular shape, electrostatics, donor/acceptor properties and polarizability as a standard series of 54 floating point numbers.

1.17 Maxima and minima of the local properties

Jakobi *et al.* [35] have described the calculation and use of the most significant maxima and minima of the local properties on the surface of the molecule. These points were used in the ParaFrag procedure to detect scaffold hops with high similarity and can be viewed as pharmacophore points.

1.18 Atom-centered descriptors

Hennemann *et al.* [36] have used atom-centered quantities calculated by ParaSurf™ as descriptors in order to calculate the strengths of hydrogen bonds [36a] and for chemical reactivity models [36b]. These descriptors (based on conventional solvent-accessible surface areas [33] using Bondi van der Waals radii [37] and a default solvent radius of 1.4 Å), C-H bond orders for hydrogen atoms, the constitution of the localized lone-pair orbitals on nitrogen atoms and the π -charges of carbon atoms in conjugated π -systems. These descriptors are now output by ParaSurf'12™.

1.19 Fragment analysis

ParaSurf'12™ can divide the input molecule into fragments (which must be defined in the input SDF file) and perform a full surface analysis for each fragment. This option and its output will be described in detail below.



2 PROGRAM OPTIONS

2.1 Command-line options

ParaSurf™ program options are given as command-line arguments. Arguments are separated by blanks, so that no single argument may contain a blank character. Arguments may be written in any combination of upper and lower case. The options are:

Table 4 ParaSurf™ command-line options

<name>		Base name for the input file (must be the first argument). <name> is not required if the first argument is -version (see below)
Using this option, the input file is assumed to be		<name>.vwr
if a file with this name exists.		
Otherwise an SDF file will be used as input in the order given.		<name>_v.sdf <name>.sdf
If neither of these files are found, the program will use an .sdf file written by the Cepos InSilico version of Mopac 6.		
These files are called		<name>_m.sdf
The output files are		<name>_p.out <name>_p.sdf <name>.psf (optional) <name>.asd (optional) <name>_p.vmp (optional)
surf=	wrap cube	Shrink-wrap surface (default) Marching-cube surface
contour=	isoden solvex solass	The surface is defined by the electron density A solvent-excluded surface is used. A solvent-accessible surface is used.
fit=	sphh isod none	Spherical-harmonic fitting (default for surf=wrap) Smooth to preset isodensity value (default for surf=cube) No fitting



iso=	n.nn	Isodensity value set to n.nn e\AA^{-3} (default for shrink-wrap surface = 0.0005 ; default for marching-cube surface = 0.007 ; minimum possible value = 0.00001)
rsol=	n.nn	A solvent-probe radius of n.nn Å is used for calculating the solvent-excluded or solvent-accessible surface (default= 1.0 , allowed range is from 0.0 to 2.0 Å)
mesh=	n.nn	The mesh size used to triangulate the surface is set to n.nn Å (default value = 0.2 Å , allowed range is from 0.1 to 1.0 Å)
estat=	naopc multi newmp	Use NAO-PC electrostatics Use multipole electrostatics (gives ParaSurf'11™ electrostatics with the parasurf11 keyword, otherwise ParaSurf'12™) Use ParaSurf'12™ multipole electrostatics (default)
psf=	on off	Write .psf surface file Do not write .psf surface file (default)
asd=	on off	Write anonymous SD (.asd) file Do not write .asd file (default)
vmp=	on off mep iel eal pol har eng anr fnm sha <MOD>	Write .vmp file for debugging. Map the MEP onto the surface Do not write .vmp file (default) Write .vmp file for debugging. Map the MEP onto the surface Write .vmp file for debugging. Map IE _L onto the surface Write .vmp file for debugging. Map EA _L onto the surface Write .vmp file for debugging. Map α_L onto the surface Write .vmp file for debugging. Map η_L onto the surface Write .vmp file for debugging. Map χ_L onto the surface Write .vmp file for debugging. Map the number of the atom assigned to the surface element onto the surface Write .vmp file for debugging. Map F _N onto the surface Write .vmp file for debugging. Map the Shannon entropy onto the surface Write .vmp file for debugging. Map the local property with the three-character designator <MOD> defined in the SIM file onto the surface
vmpfrag=	on off all	Equivalent to vmp= , but writes separate .vmp files for each fragment with only its atoms and the MEP projected onto the fragment surface. The files are named <filename><fragmentname>.vmp , where <fragmentname> is the name assigned to the fragment in the input SDF file. No fragment .vmp files will be written. As for on , except that the atoms for the entire molecule are written to the .vmp files with the surface for the fragment only.



grid=	<filename>	Read the Cartesian coordinates at which to calculate a grid of ($\log_{10}(\rho)$, MEP, IE _L , EA _L , α_L , χ_L , η_L and their first derivatives in x, y and z-directions). See Section 3.9.1 .
	auto	ParaSurf™ calculates an automatic grid that excludes areas closer than 0.5 Å to the nuclei (see Section 3.9.2)
	vdw	ParaSurf™ calculates an automatic grid that excludes areas closer than the corresponding van der Waals radius to the nuclei
	box	ParaSurf™ calculates an automatic grid including all points regardless of their proximity to nuclei
	surf	The properties of the surface points are written to the .psf file
lattice=	n.nn	Sets the lattice spacing for the grid=auto , vdw or box option (see Section 3.9.2)
sim=	<filename>	One or more surface-integral models will be read from the file <filename>.sim in the ParaSurf™ root directory. <filename> can be upper or lower case or any mixture but must be exactly three characters long.
center=	on	The atomic and surface coordinates in the .psf output file will be centered for calculations that use spherical-harmonic fitting. Note that this means that the atomic coordinates in the SDF-output file (which are the input coordinates) will be different to those in the PSF-output file. This option is default.
or		
centre=	off	The atomic and surface coordinates in the .psf output file will not be centered and will correspond to the input coordinates and those in the SDF-output file.
shannon	=<filename>	Requests that Shannon entropies (both internal and external) be calculated. If no statistics file <filename> is given, the default file (bins.txt in the ParaSurf™ Root directory) will be used. If a statistics file is given that either does not exist, contains errors or is derived from ParaSurf™ runs using different options to the current one, only the internal Shannon entropy is calculated.
autocorr	=<filename>	Requests that the surface autocorrelation functions be calculated and written to the output .sdf file. <filename> must be a ParaSurf™ output .sdf file that contains the autocorrelation functions. In this case, similarities between the two molecules will be calculated and printed (see also aclist=).
table=	<filename>	An ASCII table of the ParaSurf™ descriptors will be written to the file <filename> . If <filename> exists, the values for the current molecule will be appended to the existing table, otherwise the file will be created.
aclist=	<filename>	An ASCII table of the calculated autocorrelations will be written to the file <filename> . A total of 448 variables (7 properties in 64 bins each) are written for each molecule.
aslist=	<filename>	An ASCII table of the calculated autocorrelation similarities will be written to the file <filename> . If <filename> exists, the values for the current molecule will be appended to the existing table, otherwise the file will be created.



riflist= <filename>	An ASCII table of the calculated a standard rotationally invariant fingerprint (RIF) will be written to the file <filename> . If <filename> exists, the values for the current molecule will be appended to the existing table, otherwise the file will be created.
translate =n.nn	Requests that ParaSurf™ performs low-resolution spherical-harmonic fits using translated centers at $(+n.nn, 0, 0)$, $(-n.nn, 0, 0)$, $(0, +n.nn, 0)$, $(0, -n.nn, 0)$, $(0, 0, +n.nn)$ and $(0, 0, -n.nn)$ relative to the original center. The default value of n.nn is 0.5 Å. This value is obtained if translate is used alone. The maximum value of n.nn allowed is 1.0 Å.
translate2 =n.nn	Requests that ParaSurf™ performs a more detailed translation scan with low-resolution spherical-harmonic fits using translated centers at $(+n.nn, 0, 0)$, $(+2n.nn, 0, 0)$, $(-n.nn, 0, 0)$, $(-2n.nn, 0, 0)$, $(0, +n.nn, 0)$, $(0, +2n.nn, 0)$, $(0, -n.nn, 0)$, $(0, -2n.nn, 0)$, $(0, 0, +n.nn)$, $(0, 0, +2n.nn)$, $(0, 0, -n.nn)$ and $(0, 0, -2n.nn)$ relative to the original center. The default value of n.nn is 0.25 Å. This value is obtained if translate2 is used alone. The maximum value of n.nn allowed is 0.5 Å.
fragments	Perform a fragment analysis. The fragments must be defined in the input SDF file
desfile= <filename>	Write the binned SIM descriptors to the file <filename> . If <filename> exists, the values for the current molecule will be appended to the existing table, otherwise the file will be created. The descriptors are written as a comma-separated table with headers. Note that desmodel must also be defined.
desmodel= <code>	The bin definitions for the model denoted by <code> will be used to calculate the descriptors for the table of binned SIM descriptors. The possible values of <code> and their definitions are given in Table 2 .
-version	Must be the first argument. Requests that ParaSurf™ prints the version number to the standard output channel and then stops without performing a calculation.
eal09	Do not use the selection procedure for virtual orbitals [11] when calculating the local electron affinity. This option provides continuity with earlier versions of ParaSurf™
parasurf11	Backwards compatibility option: electron densities, local properties and electrostatic potential and field are calculated using the algorithms from ParaSurf'11™
precise	More precise output of the local properties in grid calculations
locpol= aniso	Use the local polarizability calculated from anisotropic atomic polarizability tensors (default)
 old	Use isotropic atomic polarizabilities to calculate the local polarizability (implied by the parasurf11 option)
no_derivatives	Do not calculate the first derivatives of the local properties (default is to calculate the derivatives)



Examples:

```
parasurf test surf=wrap fit=sphh iso=0.03 psf=on estat=naopc
```

Use the input file **test_v.sdf**, **test.sdf** or **test_m.sdf** to calculate a shrink-wrap surface with an isodensity value of 0.03 e \AA^{-3} , perform a spherical-harmonic fit, use NAO-PC electrostatics and write the spherical-harmonic coefficients to **test_P.sdf** and the entire surface to **test_P.psf**.

```
parasurf test surf=cube fit=none
```

Use the file **test_v.sdf**, **test.sdf** or **test_m.sdf** as input to perform a marching-cube surface determination without fitting and to calculate the descriptor set.



2.2 Options defined in the input SDF-file

2.2.1 Defining the center for spherical-harmonic fits

The automatic determination of the molecular center for spherical-harmonic fitting can be overridden by adding a field to the Input (usually VAMP) SDF-file with the tag:

<SPHH_CENTER>

The center can be defined using Cartesian coordinates using an input line (immediately after the **SPHH_CENTER** tag) of the format:

Cartesian x.xxy.yyz.zz

where **x.xx**, **y.yy** and **z.zz** are the x, y, and z-coordinates, respectively. The capitalization of “**Cartesian**” is required.

Alternatively, a list of atoms can be given using the format

Atoms n1 n2 n3 n4 n5 n6 ...

where **n1** etc. are the numbers of the atoms to be used to calculate the center of gravity. The capitalization of “**Atoms**” is required and the list of atoms is limited to one line.



2.2.2 Defining fragments

Molecular fragments can be defined in the input SDF file and fragments calculations requested using the fragments options, for instance

```
parasurf test surf=cube fragments
```

Figure 9 shows a sample <fragment> block from an SDF input file.

```
><FRAGMENTS>
Start fragment
phenyl
 3 4 5 15 16 19 25 33
End fragment

Start fragment
methoxyl
 1 2 22 23 24
End fragment

Start fragment
methoxy2
17 18 34 35 36
End fragment

Start fragment
methoxy3
20 21 37 38 39
End fragment

Start fragment
methylene
6 26 27
End fragment

Start fragment
thymine
 7 8 9 10 11 12 13 14 28 29 30 31 32
End fragment

Start fragment
everybody
 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20
 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39
End fragment

><END_FRAGMENTS>
```

Figure 9 A sample <FRAGMENTS> input block



The first line after each “**Start fragment**” line (note the upper and lower case, which are necessary) defines the name given to the fragment. This is followed by the numbers of the atoms that make up the fragment (20i4, fixed format). Note that the fragments need not be mutually exclusive. The fragment “**everybody**” in the above example, for instance is the entire molecule. The fragment-definition block begins with

><FRAGMENTS>

and ends with

><END_FRAGMENT>

tags.

Figure 10 shows the input molecule and the fragments.




Figure 10 The fragments defined in the SDF input example

In a FRAGMENTS run, ParaSurf™ first performs a calculation for the entire molecule and then analyses the molecular surface according to the standard ParaSurf™ technique used to assign surface triangles to individual atoms. The output for the phenyl fragment is shown in Figure 11. A similar output section is printed for each fragment. The results and the descriptors for each fragment are taken from the surface for the whole molecule and therefore refer to the fragment (both its electronic properties and the area of its surface) within the context of the molecule itself. The coordinates given for the maxima and minima of the local properties refer to the input geometry of the entire molecule.



```
<> Results for fragment number 1 : phenyl
Surface area      :        47.21 Angstrom**2
Fragment charge   :       -0.01

          MEP      IEL      EAL      HARD      ENEG      F (N)      POL
Mean     :      -6.0      402.5     -67.6     235.1     167.5      -4.0      0.76
Mean +ve:      10.9           0.0           8.6
Mean -ve:     -13.0     -67.6      -8.7
Maximum   :      18.9      525.3     -19.9     303.2     222.2      16.9      1.4
Minimum   :     -53.1      337.6    -103.9     191.6     142.3     -42.4      0.33
Variance:      200.3      919.4      562.5     594.8     146.2      85.0      0.54E-01
Var.+ve:      29.8           0.0           21.9
Var.-ve:      103.2      562.5           28.5
Balance   :      0.174           0.000           0.246
Skew     :      -0.4           0.7           0.2      -0.1           1.1           1.6      -0.41
Kurtosis:      0.4           0.6           -1.3      -1.0           1.7           -0.2      -0.72

MEP Maxima for this fragment
Number      x          y          z      MEP
4 :      1.6569      1.8397     -2.5118     15.9683
6 :     -4.3431     -1.6853     -1.1618     18.8870

IEL Maxima for this fragment
Number      x          y          z      IEL
1 :      1.6861     -2.6103     -3.4142     525.3250

EAL Maxima for this fragment
Number      x          y          z      EAL
1 :     -1.4714     -2.0603     -3.9285    -22.4779
2 :      0.0069     -1.2520     -4.1285    -19.8861
3 :     -2.5431     -0.8270     -3.5285    -27.9933
4 :      0.8569     -1.1103     -0.6285    -25.1242
6 :     -0.7431     -2.4270     -0.4785    -28.5599

POL Maxima for this fragment
Number      x          y          z      POL
1 :     -0.7431     -3.7603     -3.4785     1.3387
2 :     -0.8648     -3.8603     -3.2785     1.3698

POL Minima for this fragment
Number      x          y          z      POL
23 :     -4.1431     -1.0103     -2.5285     0.3418
33 :      1.6569      1.1897     -1.0285     0.3529

FN Maxima for this fragment
Number      x          y          z      FN
12 :      1.2569      2.0897     -1.7902     14.2146
15 :     -4.2931     -1.0603     -1.0285     15.7029
```

Figure 11 ParaSurf12™ output for the phenyl fragment defined above



The individual surfaces of the fragments are shown in Figure 12.




Figure 12 Surfaces calculated for the individual fragments, color coded according to the MEP in kcal mol^{-1} . The fragments (clockwise from the top right) are methoxy1, methoxy2, methoxy3, thymine, phenyl and methylene



3 INPUT AND OUTPUT FILES

ParaSurf™ uses the following files for input and output:

Table 5 ParaSurf™ input and output files

File	Name	Description
Input	<filename>.vwf (if available) or <filename>_v.sdf or <filename>.sdf or <filename>_m.sdf	EMPIRE™ .vWF file VAMP .SDF file output. VAMP must be run with the ALLVECT option to be able to calculate all the properties. The VAMP version used must be able to calculate AO-polarizabilities. If no VAMP .SDF file is found, ParaSurf™ defaults to a CeposMopac 6 .SDF file. It is strongly recommended to use the EF option for geometry optimizations in Mopac.
Hamiltonian	Vhamil.par	The VAMP parameters file (also found in the VAMP executable directory). This file must be copied to the ParaSurf™ executable directory.
Output	<filename>_p.out	Always written.
SD-file	<filename>_p.sdf	Always written.
ASD-file	<filename>.asd	Anonymous SD-file. Requested by the option asd=on
PSF-file	<filename>.psf	ParaSurf™ surface file. Requested by the option psf=on
VMP-file	<filename>_p.vmp	Debug file.
SIM-file	<filename>.sim	Surface-integral model definition. <filename> must have exactly three characters and the file must reside in the ParaSurf™ executable directory.
Descriptor table file	User defined	An ascii, comma-separated file that contains a line of descriptors for each molecule. This file will be created if it does not exist or an extra line will be appended if it does exist.
Binned SIM descriptor file	User defined	An ascii, comma-separated file that contains a line of the descriptors generated for the bin definitions used in the model defined by <code> in the desmodel= command-line option. A header defining the descriptors is printed as the first line.
Autocorrelation fingerprint file	User defined	An ascii, comma-separated file that contains the molecule's ID and 448 binned autocorrelation values. The file will be overwritten if it exists
Autocorrelation similarity file	User defined	An ascii, fixed format file that contains a line of seven autocorrelation similarities for each molecule. This file will be created if it does not exist or an extra line will be appended if it does exist.
RIF table file	User defined	An ascii, comma-separated file that contains a line of the standard rotationally invariant fingerprint (RIF [37]) for each molecule. This file will be created if it does not exist or an extra line will be appended if it does exist.



3.1 The VAMP .sdf file as input

VAMP .sdf files, an extension of the MDL .sdf file format, [38] are the primary communication channel between VAMP and ParaSurf™. The atomic coordinates and bond definitions are given in the MDL format as shown in Figure 13. The remaining fields are indicated by tags with the form:

<FIELD_NAME>

FIELD_NAME is a predefined text tag used to locate the relevant data within the .sdf file.

Only the important fields for a ParaSurf™ calculation will be described here:

```
1-Bromo-3,5-difluorobenzene
OMVAMP81A04250313563D 1    0.00000      0.00000      0

12 12 0 0 0 0           1 V2000
-2.6274   0.2410   0.0003 F
-1.2738   0.2410   0.0003 C
-0.5810   1.4623   0.0003 C
 0.8231   1.4389   0.0003 C
 1.5096   2.6055   0.0004 F
 1.5266   0.2198   0.0001 C
  0.8142  -0.9793   0.0001 C
  1.7431  -2.6055  -0.0004 Br
 -0.5805  -0.9840   0.0002 C
 -1.1264   2.4167  -0.0003 H
  2.6274   0.2339   0.0003 H
 -1.1515  -1.9253   0.0001 H
 1  2  1
 2  3  4
 3  4  4
 4  5  1
 4  6  4
 6  7  4
 7  8  1
 2  9  4
 7  9  4
 3 10  1
 6 11  1
 9 12  1
M  END
```

Figure 13 The headers and titles, atomic coordinates and bond definitions from a VAMP .sdf file. The format follows the MDL definition [26]

<HAMILTONIAN>

The Hamiltonian field defines the semiempirical Hamiltonian (model and parameters) used for the calculation. The Hamiltonian must be defined for ParaSurf™ to be able to calculate the electrostatics and the local polarizabilities. NAO-PC electrostatics and the local polarizability are not available for all methods. Quite generally, the multipole electrostatics model is to be preferred over the NAO-PC model, which can only be used if the VAMP .sdf file contains a block with the tag:

<NAO-PC>

NAO-PCs cannot be calculated for methods with *d*-orbitals. The local polarizability calculation has not yet been extended to these methods, but will be in a future release.



The following table gives an overview of the methods and their limitations:

Table 6 Hamiltonians and the available electrostatic and polarizability models.

Hamiltonian	Reference	Electrostatics		Local Polarizability
		NAO-PC	Multipole	
MNDO	[20a]	YES	YES	YES
AM1	[22]	YES	YES	YES
PM3	[23]	YES	YES	YES
MNDO/c	[39]	YES	YES	NO
MNDO/d	[20b]	NO	YES	NO
AM1*	[40]	NO	YES	NO

<VAMPBASICS>

The VAMPBASICS block contains the following quantities (FORTRAN format 6f13.6):

Heat of Formation	kcal mol ⁻¹
HOMO energy	eV
LUMO energy	eV
Dipole moment	
x-component	Debye
y-component	Debye
z-component	Debye

<TOTAL COULSON CHARGE>

The total charge of the molecule.

<DENSITY MATRIX ELEMENTS>

The DENSITY MATRIX ELEMENTS block contains the one-atom blocks of the density matrix for the non-hydrogen atoms. For an *sp*-atom, there are ten elements, for an *spd*-atom 45. The squares of the diagonal elements for hydrogen atoms are included in the <CHARGE ON HYDROGENS> block that follows the density matrix. The density-matrix elements are used in ParaSurf™ to calculate the local properties and are essential.

<ORBITAL VECTORS>

The ORBITAL VECTORS block contains the MO-eigenvectors and related information and is essential for calculating the local properties. VAMP must be run with the keyword **ALLVECT** in order to write all the MO vectors to the SDF file.

The entire SDF input file is echoed to the <filename>.p.sdf output file and the properties calculated by ParaSurf™ are added in additional blocks at the end.



3.1.1 Multi-structure SD-files

ParaSurf™ can read SD-files containing more than one molecule (e.g. those produced by the VAMP-QSAR model engine) and process them in one run. The command-line arguments apply to each molecule in the SD-file and the same semiempirical Hamiltonian must be used for each molecule or an error message will be printed and the program terminated.

As part of this enhancement, ParaSurf™ can use SD-files that do not contain the one-atom blocks of the density matrix explicitly. Thus, SD-files that only contain the molecular-orbital Eigenvectors and Eigenvalues give full ParaSurf™ functionality within the previous restrictions that:

- Polarizabilities are not yet available for Hamiltonians that use *d*-orbitals (MNDO/d and AM1*).
- NAO-PC electrostatics are only available if the NAO-PCs are present in the SD-file. Multipole electrostatics are available for all Hamiltonians.

The output SD-file written by ParaSurf™ also contains multiple molecules as in the input file. Other ParaSurf™ output files (`.asd`, `.vmp` etc.) are also concatenated.

Multiple SD-files can be used with a SIM file exactly as single molecules.

3.2 The Cepos Mopac 6 .sdf file as input

Cepos Mopac 6 writes an `.sdf` file containing the above blocks with the exception that the MOPACBASICS block replaces VAMPBASICS. No additional keywords are required to request the correct `.sdf` output for ParaSurf™.

3.3 The Vhamil.par file

The file `Vhamil.par` is used by VAMP to define the available Hamiltonians and elements and supply the parameters. This file is also used by ParaSurf™ for the same purpose. A `Vhamil.par` file for standard Hamiltonians and elements is supplied with the ParaSurf™ program. In order to be sure that all Hamiltonians and elements available to VAMP can also be handled by ParaSurf™, however, the `Vhamil.par` file from the VAMP executable directory should be copied into the ParaSurf™ executable directory. The `Vhamil.par` file is not necessary if an EMPIRE™ `.vwf` file is used as input.



3.4 The EMPIRE™ binary wavefunction file (.vwf)

The binary wavefunction file contains all the information necessary to process the results of the EMPIRE™ calculation further. It is the new primary input file format for ParaSurf™ and does not require a **Vhamil.par** file to be present. Its contents are:

Table 7 Content of the EMPIRE™ binary wavefunction file (.vwf)

Definition	Type
First comment line from the input	character (len=80)
Title line from the input	character (len=80)
Hamiltonian	character (len=6) "AM1 ", "AM1* ", "MNDO ", "MNDO/c", "MNDO/d", "PM3 " or "PM6 "
Formalism	character (len=3) "RHF"
Number of atoms	integer
Number of orbitals (Norbs)	integer
Number of doubly occupied orbitals	integer
Number of singly occupied orbitals	integer = 0
Charge on the molecule	integer
Heat of formation in kcal mol ⁻¹	double precision
Energy of the HOMO (eV)	double precision
Energy of the LUMO (eV)	double precision
x, y and z-components of the dipole moment (Debye)	double precision (1:3)
For each atom:	
	Atomic number
	Cartesian coordinates
	Number of atomic orbitals
	Principal quantum number (s and p)
	Principal quantum number (d)
	Slater exponents (ζ_s , ζ_p , ζ_d)
	Overlap integrals (β_s , β_p , β_d)
	Multipole parameters (dd and qq)
End atoms	
Eigenvalues (eV)	double precision (1:Norbs)
Eigenvectors	double precision (1:Norbs ²)




3.5 The ParaSurf™ output file

The ParaSurf™ output file provides the user with information about the calculation and the results. It is, however, not intended as the primary means of communication between ParaSurf™ and other programs. Thus, the essential information contained in the output file is also available from the ParaSurf™ output **.sdf** file.

3.5.1 For a spherical-harmonic surface

Figure 14 shows the output for a calculation using the options **surf=wrap fit=sphh translate** for trimethoprim, 1.



ParaSurf™ output for trimethoprim, 1, using a spherical-harmonic surface

```
<> ParaSurf'12, Revision 410
<> Copyright (c) 2006-2012 Friedrich-Alexander-Universitaet Erlangen-Nuernberg
     and Cepos InSilico Ltd. All rights reserved.

<> Input = trimethoprim.sdf

<<>> Molecule      1 of      1 <<>>

<> Program options :

Using shrink-wrap isocontour surface
Fitting surface to spherical harmonics
Translations for spherical-harmonic fits: 1 step of      0.5000 Angstrom in each
direction.
Using an isodensity surface contour
Isodensity value =  0.5000E-03 electrons/Angstrom**3
Triangulation mesh =      0.20 Angstrom
Using multipole electrostatics

<> AM1      calculation for   Trimethoprim

<> Translated spherical-harmonic fits:
      dx      dy      dz      rmsd
    0.0000  0.0000  0.0000  0.3827
    0.5000  0.0000  0.0000  0.5516
   -0.5000  0.0000  0.0000  0.4939
```



ParaSurf™ output for trimethoprim, 1, using a spherical-harmonic surface

```
0.0000  0.5000  0.0000  0.5333
0.0000 -0.5000  0.0000  0.5173
0.0000  0.0000  0.5000  0.5406
0.0000  0.0000 -0.5000  0.4147

<> Fitting surface to spherical harmonics
<> Order(1)    RMSD

      0      2.01177719
      1      2.06264149
      2      1.57429428
      3      1.14144633
      4      0.96586545
      5      0.69034677
      6      0.60076280
      7      0.51821086
      8      0.48396719
      9      0.45955001
     10      0.42519621
     11      0.39310379
     12      0.37725481
     13      0.36600493
     14      0.34829905
     15      0.32860993

<> Spherical harmonic fit for MEP:
<> Order(1)    RMSD

      0      12.40803637
      1      12.34784414
      2      9.25211030
      3      8.29692710
      4      6.82440697
      5      5.74002976
      6      4.82285463
      7      4.20321063
      8      3.80848913
      9      3.53212800
     10      3.08074911
     11      2.62541255
     12      2.37675753
     13      2.06595225
     14      1.98683370
     15      1.84188725
     16      1.70418851
     17      1.54855928
     18      1.34322707
     19      1.21349689
     20      1.08035065

<> Spherical harmonic fit for IE(1):
<> Order(1)    RMSD

      0      44.57878887
      1      39.49221723
      2      37.79258047
      3      36.00326805
      4      32.26724224
      5      29.12695030
      6      26.30443933
      7      25.27443867
      8      23.78306670
      9      22.00750560
     10      21.22324848
     11      20.31770921
     12      19.17042684
     13      17.99969469
     14      17.10345128
     15      16.83692340
     16      16.18125074
```



ParaSurf™ output for trimethoprim, 1, using a spherical-harmonic surface

```
17      14.74040808
18      13.59009122
19      12.96531563
20      12.96531563

<> Spherical harmonic fit for EA(l):
<> Order(1)   RMSD

0      14.76660122
1      14.57921630
2      14.82522427
3      11.54502778
4      10.82050211
5      9.79342370
6      9.76520958
7      9.37069356
8      8.72479764
9      8.01299396
10     7.52757624
11     7.18259970
12     6.90528697
13     5.87616823
14     5.39515470
15     4.92869560
16     4.68556059
17     4.48184805
18     4.20875146
19     4.02983055
20     3.88447733

<> Spherical harmonic fit for Field(N):
<> Order(1)   RMSD

0      10.23375986
1      10.22721474
2      9.38433421
3      9.05353285
4      8.17889165
5      7.54963618
6      6.97197468
7      6.65848164
8      6.39618968
9      6.09545254
10     5.46552427
11     4.86094641
12     4.46112161
13     4.12120028
14     4.03929276
15     3.83138764
16     3.55514598
17     3.21320455
18     2.77835041
19     2.50117320
20     2.28148569

<> Spherical harmonic fit for Alpha(l):
<> Order(1)   RMSD

0      0.24597908
1      0.24673395
2      0.23405408
3      0.20695856
4      0.18909827
5      0.17468336
6      0.16936390
7      0.15804017
8      0.14856557
9      0.13972827
10     0.12529394
11     0.11480278
12     0.10565425
```



ParaSurf™ output for trimethoprim, 1, using a spherical-harmonic surface

13	0.09908593
14	0.09602514
15	0.09019338
16	0.08240006
17	0.07510580
18	0.06885870
19	0.06580610
20	0.06287666
<> Property ranges:	
Density	: 0.1004E-03 to 0.1836E-02
IE(1)	: 325.65 to 554.64
EA(1)	: -112.61 to -29.37
MEP	: -50.22 to 19.02
Alpha(1)	: 0.2167 to 1.5089
Field(N)	: -52.89 to 19.81
<> Descriptors :	
Dipole moment	: 1.2467 Debye
Dipolar density	: 0.2445E-02 Debye.Angstrom**-3
Molecular pol.	: 128.5408 Angstrom**3
Molecular weight	: 290.32
Globularity	: 0.7559
Total surface area	: 408.28 Angstrom**2
Molecular volume	: 509.81 Angstrom**3
Most positive MEP	: 19.02 kcal/mol
Most negative MEP	: -50.22 kcal/mol
Mean +ve MEP	: 6.28 kcal/mol
Mean -ve MEP	: -11.89 kcal/mol
Mean MEP	: -2.37 kcal/mol
MEP range	: 69.24 kcal/mol
MEP +ve Variance	: 15.50 (kcal/mol)**2
MEP -ve Variance	: 133.21 (kcal/mol)**2
MEP total variance	: 148.71 (kcal/mol)**2
MEP balance parameter	: 0.0934
MEP balance*variance	: 13.8883 kcal/mol
MEP skewness	: -1.3157
MEP kurtosis	: 1.5929
Integral MEP	: -823.474 kcal.Angstrom**2/mol
Maximum IE(1)	: 554.64 kcal/mol
Minimum IE(1)	: 325.65 kcal/mol
Mean IE(1)	: 422.44 kcal/mol
IE(1) range	: 228.99 kcal/mol
IE(1) variance	: 1947.25 (kcal/mol)**2
IE(1) skewness	: 0.6846
IE(1) kurtosis	: -0.3181
Integral IE(1)	: 7450.37 eV.Angstrom**2
Maximum EA(1)	: -29.37 kcal/mol
Minimum EA(1)	: -112.61 kcal/mol
Mean +ve EA(1)	: 0.00 kcal/mol
Mean -ve EA(1)	: -93.26 kcal/mol
Mean EA(1)	: -93.26 kcal/mol
EA(1) range	: 83.24 kcal/mol
EA(1) +ve variance	: 0.00 (kcal/mol)**2
EA(1) -ve variance	: 203.49 (kcal/mol)**2
EA(1) total variance	: 203.49 (kcal/mol)**2
EA(1) skewness	: 1.5615
EA(1) kurtosis	: 3.0484
Integral EA(1)	: -1647.42 eV.Angstrom**2
EA(1) balance param.	: 0.0000
Fraction pos. EA(1)	: 0.0000 (= 0.00 Angstrom**2)
Max. local Eneg.	: 234.22 kcal/mol
Min. local Eneg.	: 113.00 kcal/mol
Mean local Eneg.	: 164.59 kcal/mol
Local Eneg. range	: 121.21 kcal/mol
Local Eneg. variance	: 453.76 (kcal/mol)**2



ParaSurf™ output for trimethoprim, 1, using a spherical-harmonic surface

```

Local Eneg. skewness :      0.66
Local Eneg. kurtosis :     0.01
Integral local Eneg. : 2901.47 eV.Angstrom**2

Max. local hardness :    321.14 kcal/mol
Min. local hardness :   187.77 kcal/mol
Mean local hardness : 257.85 kcal/mol
Local hard. range :    133.37 kcal/mol
Local hard. variance : 621.61 (kcal/mol)**2
Local hard. skewness :  0.39
Local hard. kurtosis : -0.39
Integral local Hard. : 4548.90 eV.Angstrom**2

Maximum alpha(1) :      1.509 Angstrom**3
Minimum alpha(1) :     0.2167 Angstrom**3
Mean alpha(1) :        0.4780 Angstrom**3
Alpha(1) range :      1.292 Angstrom**3
Variance in alpha(1) : 0.5893E-01 Angstrom**6
Alpha(1) skewness :    1.5346
Alpha(1) kurtosis :    1.5720
Integral Alpha(1) :   194.797 Angstrom**5

Maximum field normal : 19.81 kcal/mol.Angstrom
Minimum field normal : -52.89 kcal/mol.Angstrom
Mean field :          -0.27 kcal/mol.Angstrom
Field range :         72.70 kcal/mol.Angstrom
Total field variance : 104.50 (kcal/mol.Angstrom)**2
+ve field variance : 11.74 (kcal/mol.Angstrom)**2
-ve field variance : 116.79 (kcal/mol.Angstrom)**2
Field balance param. : 0.08
Field skew :           2.46
Field kurtosis :       5.021
Integral F(N) :        44.29 kcal.Angstrom/mol
Integral F(N +ve) :   1456. kcal.Angstrom/mol
Integral F(N -ve) :  -1411. kcal.Angstrom/mol
Integral |F(N)| :    2867. kcal.Angstrom/mol

<> Spherical-Harmonic Hybridization:

Shape hybrids :
16.009628  1.373167  3.546672  2.694125  1.350668  1.644624
0.762716  0.658342  0.375899  0.356134  0.403193  0.301343
0.190579  0.199740  0.230850  0.204322

MEP hybrids :
7.934381  4.585824  29.916318  11.316554  14.111406  11.572654
10.218143 8.395453  5.647876  4.106887  4.808912  4.310994
3.286470  3.516688  1.782163  2.046608  1.478736  1.563289
1.807359  1.409069  1.370528

IE(1) hybrids :
1475.0734  69.4741  57.6041  47.8482  44.9670  44.0421
43.8721  28.8369  37.2336  34.3592  20.0558  16.1335
20.3418  18.4038  18.1736  13.1804  13.3178  17.9975
17.9273  16.0483  0.0000

EA(1) hybrids :
317.0698  7.7090  18.9862  33.3973  15.7480  17.4261
11.2668  9.8258  11.5616  10.4208  9.5707  8.2159
6.5635  9.7503  6.2945  7.1239  4.4548  4.3692
4.8386  4.2292  3.5039

Alpha(1) hybrids :
1.83506449 0.07660952 0.42325808 0.37116637 0.26168092 0.25387096
0.19306697 0.22512754 0.17314967 0.16214546 0.15522714 0.12040044
0.12344010 0.10628415 0.08308407 0.08170594 0.07634731 0.07232668
0.07259004 0.05268140 0.04875321

Field(N) hybrids :
2.6299  2.0991  15.8335  8.7229  12.1467  10.8157
9.8725  9.3441  7.0443  5.6697  7.0721  6.5543

```



ParaSurf™ output for trimethoprim, 1, using a spherical-harmonic surface

6.2485	6.0541	3.3904	3.6946	3.2560	3.7721
4.2037	3.4602	2.9100			

<> Standard rotationally invariant fingerprint: (L. Mavridis, B. D. Hudson and D. W. Ritchie, J. Chem. Inf. Model., 2007, 47, 1787-1796.)

4.00120	1.17182	1.88326	1.64138	1.16218
1.28243	0.873336	2.81680	2.14145	5.46958
3.36401	3.75652	3.40186	3.19658	2.89749
2.37653	2.02655	2.19292	2.07629	1.81286
38.4067	8.33511	7.58974	6.91724	6.70575
6.63643	6.62360	17.8065	2.77652	4.35732
5.77904	3.96837	4.17446	3.35661	1.35465
0.276784	0.650583	0.609234	0.511548	0.503856
0.439394	1.62171	1.44884	3.97914	2.95346
3.48521	3.28872	3.14205	3.05681	2.65411
2.38110	2.65934	2.56013	2.49969	

<> Atomic surface properties:

Atom	Area	MEP		IE(1)		EA(1)		mean pol.	Field(N)		Energ(L)		Hard(L)		
		max	min	max	min	max	min		max	min	max	min	max	min	
C	1	1.190	-4.99	-24.60	474.40	440.53	-84.14	-97.49	0.695	-0.83	-19.62	195.13	171.52	279.27	268.74
O	2	3.298	-15.21	-48.75	483.72	395.45	-68.96	-88.24	1.013	-10.63	-35.96	204.79	153.60	278.92	238.20
C	3	5.583	2.76	-42.26	473.24	343.23	-31.92	-102.63	0.845	3.14	-23.16	199.07	137.84	274.57	192.42
C	4	2.301	-1.09	-11.01	420.74	358.07	-47.79	-88.79	0.765	0.11	-7.46	167.86	147.78	252.89	202.93
C	5	1.315	-2.83	-11.04	443.62	381.62	-69.72	-93.99	0.780	-2.54	-6.76	174.81	152.40	268.81	226.23
C	6	0.000													
C	7	1.851	-1.56	-11.81	420.70	353.35	-50.94	-93.17	0.866	-0.53	-7.17	169.93	141.70	250.77	207.08
C	8	6.600	0.72	-22.08	435.21	362.23	-35.07	-99.15	0.779	2.08	-15.47	176.51	150.00	258.95	202.94
N	9	7.657	-9.91	-40.75	443.85	328.33	-82.12	-109.10	1.186	-10.05	-52.89	175.00	113.00	269.71	214.57
C	10	10.568	5.91	-23.31	493.46	388.30	-37.01	-96.70	0.791	4.70	-15.87	208.30	154.49	288.55	213.45
N	11	1.649	-14.33	-23.08	511.29	446.71	-63.02	-82.87	0.793	-16.61	-22.27	216.19	190.10	295.11	254.86
N	12	6.592	-9.43	-39.67	444.68	325.65	-80.18	-106.59	1.031	-10.86	-51.95	172.11	114.98	273.94	206.65
C	13	7.498	4.98	-25.74	511.74	366.79	-34.15	-96.77	0.813	6.63	-21.17	215.43	152.07	296.31	201.46
N	14	1.811	-14.21	-22.59	489.48	443.66	-66.69	-87.80	0.801	-18.53	-29.45	206.02	184.52	285.56	256.30
C	15	4.876	-0.83	-11.93	434.24	358.03	-52.14	-100.50	0.747	2.70	-9.39	173.03	145.57	261.21	205.09
C	16	5.487	-2.05	-37.60	476.55	346.16	-29.37	-99.54	0.821	2.98	-22.97	200.45	139.16	276.10	187.77
O	17	0.967	-23.73	-38.37	475.37	409.71	-71.36	-88.47	0.959	-11.16	-33.69	198.69	165.37	276.68	241.01
C	18	0.868	9.69	-22.92	505.17	434.80	-83.99	-98.88	0.690	5.54	-12.51	208.01	167.96	297.16	266.84
C	19	5.387	-1.43	-46.05	485.13	325.82	-36.16	-103.21	1.007	-5.13	-29.69	201.97	133.24	286.16	189.70
O	20	4.532	-34.23	-50.22	477.62	372.36	-73.80	-95.94	1.086	-18.29	-43.77	201.91	143.17	279.48	228.85
C	21	1.026	-20.40	-37.71	465.20	425.64	-93.36	-109.78	0.771	-4.77	-21.36	184.55	157.93	280.68	267.71
H	22	24.370	9.65	-29.35	508.71	383.58	-87.48	-97.43	0.402	6.63	-16.30	209.44	144.98	299.27	238.60
H	23	17.654	12.04	-11.12	503.85	382.54	-76.95	-97.30	0.370	7.51	-8.90	207.29	145.64	296.55	236.85
H	24	18.559	11.71	-14.40	502.99	382.80	-77.32	-96.21	0.366	6.31	-3.91	206.62	145.37	296.38	237.44
H	25	6.426	12.26	0.38	450.48	398.83	-84.46	-99.09	0.375	9.18	-4.08	178.56	152.24	271.92	243.04
H	26	14.980	8.87	-4.47	463.19	374.80	-88.84	-102.54	0.381	6.24	-3.25	182.47	139.03	281.34	235.77
H	27	14.448	8.05	-15.74	466.07	374.02	-85.09	-102.86	0.384	5.61	-6.12	190.49	137.76	279.96	236.26
H	28	18.509	9.52	-25.29	438.41	392.07	-70.47	-102.76	0.407	8.09	-22.53	174.36	149.49	270.08	233.30
H	29	23.099	18.93	-25.76	550.91	428.54	-78.35	-111.77	0.334	19.12	-24.36	232.83	159.04	321.14	269.51
H	30	22.784	19.02	-25.71	549.21	416.66	-78.19	-111.79	0.316	19.81	-23.60	234.22	154.05	320.21	262.61
H	31	22.910	15.18	-25.53	554.64	420.49	-77.17	-112.61	0.314	13.89	-26.30	234.06	155.95	320.58	264.54
H	32	8.916	14.53	-11.92	552.74	397.85	-78.64	-104.83	0.357	14.60	-19.29	232.60	149.52	320.13	248.33
H	33	6.289	10.08	-4.74	441.04	384.41	-73.12	-98.75	0.443	8.53	-6.17	172.27	147.83	268.77	228.76
H	34	23.001	9.47	-23.20	498.69	384.37	-87.93	-99.94	0.374	6.36	-15.55	204.62	145.09	294.07	238.62
H	35	18.462	9.75	-15.61	500.50	380.61	-73.19	-96.24	0.380	6.52	-6.87	205.69	145.35	294.81	231.10
H	36	18.392	9.64	-6.07	502.41	382.26	-80.84	-97.20	0.382	8.17	-12.02	206.76	145.20	295.65	237.06
H	37	24.577	3.25	-31.68	487.62	370.69	-99.58	-109.64	0.394	3.95	-22.89	192.74	133.11	294.88	237.57
H	38	17.255	2.55	-34.53	495.67	367.07	-87.27	-109.17	0.387	2.36	-21.41	196.52	130.24	299.15	236.77
H	39	21.193	2.66	-35.53	490.81	359.03	-87.36	-108.60	0.376	2.69	-25.26	194.23	130.19	296.58	226.39

Total 402.881

<> Stationary points on the molecular surface (A. Jakobi, H. Mauser and T. Clark, J. Mol. Model., 2008, 14, 547-558)

x	y	z	value
<> 9 MEP Maxima :			
-2.0943	-1.9103	1.9958	14.48
1.9308	2.6575	-1.9070	10.08
4.2004	2.2889	-1.9835	9.694
1.4638	5.7589	5.4770	12.75
-1.4292	-1.7902	3.3075	15.18
-0.1036	3.5080	-2.4658	9.524



ParaSurf™ output for trimethoprim, 1, using a spherical-harmonic surface

-4.5244	-1.3404	-0.6240	12.26
1.9409	3.3233	5.3060	17.29
2.3347	3.9929	4.8681	19.02
<> 5 MEP Minima :			
0.6146	5.8476	1.0368	-40.75
0.0000	-4.3478	-2.5240	-50.22
-1.4714	2.3729	4.3876	-39.67
0.8011	-3.6508	-3.4297	-45.17
2.6850	-2.2208	-2.9412	-38.37
<> 7 IEL Maxima :			
-3.0425	-1.2046	2.7457	554.6
0.2652	-0.8162	2.3579	538.2
-0.1428	-1.3587	2.3663	539.8
-1.9644	-1.8860	2.7447	546.7
-0.9455	-1.8557	2.5130	543.5
2.1930	5.5001	5.4578	550.9
1.0590	5.7161	5.3691	550.9
<> 4 IEL Minima :			
-1.3070	2.2637	4.5275	325.7
0.2228	-2.1196	-3.6915	325.8
-0.2634	1.6256	4.6963	332.5
0.9079	5.7325	1.0276	328.3
<> 4 EAL Maxima :			
-2.9083	2.6186	2.2719	-34.15
-2.2332	4.3872	0.4313	-35.07
-1.0955	-2.0487	-3.3586	-31.92
0.7486	-1.0304	-3.6793	-29.37
<> 8 EAL Minima :			
0.2957	-5.5123	-1.4812	-109.8
1.1122	-5.2323	-3.1054	-107.1
-0.9624	-0.1287	4.9079	-112.6
2.6858	-2.4183	0.9711	-106.7
3.1809	-3.6014	-3.0616	-109.3
4.1893	-2.5658	0.2149	-106.6
0.5536	2.7836	5.8582	-111.8
1.8617	6.2852	2.3982	-111.8
<> 1 Alpha(1) Maxima :			
0.9079	5.7325	1.0276	1.509
<> 43 Alpha(1) Minima :			
0.1941	-0.5973	3.5616	0.2183
-2.6409	-5.9315	0.0000	0.3151
0.0000	-5.4569	0.0000	0.3507
-3.8091	2.1992	-1.6009	0.3088
-3.2197	2.3393	-2.2977	0.3049
4.7632	-0.6017	-4.0286	0.3010
2.0757	1.6767	0.7160	0.3543
-1.7867	4.6716	-1.3420	0.2991
-3.8319	1.4763	1.1018	0.3538
-3.3602	-0.9081	1.6253	0.2685
0.0000	-1.1114	3.2106	0.2167
-1.8684	-5.7503	-2.2128	0.3172
2.5578	-5.0048	1.5081	0.3072
1.3439	3.4788	-1.7414	0.3159
1.3551	2.3352	-2.7118	0.3274
-4.8888	-2.1766	0.0000	0.3256
3.1679	0.2910	-4.5679	0.2972
2.2360	1.3732	0.8281	0.3515
-2.4257	0.7858	4.4899	0.2262
-2.5222	-3.6775	1.4074	0.3008
0.1013	-3.7287	1.1772	0.3636
-1.7602	-5.9235	-1.3715	0.3161
-5.0212	-3.6481	0.5445	0.3369
-2.4804	-5.5710	-2.8512	0.3166
4.0476	-0.2121	0.7176	0.2939
2.4151	5.0624	5.1803	0.2208
3.0686	2.6189	-0.8974	0.3557
-0.4681	4.4656	5.9016	0.2285
4.6449	-3.9692	0.2672	0.3134



ParaSurf™ output for trimethoprim, 1, using a spherical-harmonic surface				
4.4728	-3.4303	-1.7844	0.3240	
-4.8828	-1.8743	-1.9115	0.3157	
-3.9258	-0.7208	-2.0799	0.3162	
-3.3468	-3.7159	-3.8577	0.3006	
3.5219	-6.1001	-0.6180	0.3331	
3.0798	-4.7429	-3.2785	0.3043	
5.4843	-1.0145	-0.7349	0.2994	
5.7718	1.0663	-2.4400	0.3606	
1.2318	-3.0701	1.4574	0.3178	
2.0933	5.6781	2.0628	0.2651	
2.5348	4.8420	2.6986	0.2362	
2.0840	3.3480	4.9571	0.2216	
2.6333	4.8195	3.3405	0.2301	
0.3362	6.5480	2.7231	0.2358	
<> 13 F(N) Maxima :				
-0.5388	-1.6584	2.0781	14.28	
1.1425	5.9891	5.1567	12.89	
-1.6503	-1.7170	3.4429	13.58	
0.3958	5.0472	6.0897	10.52	
2.1539	3.5125	4.9561	17.87	
-0.2643	0.4295	4.5979	11.42	
-1.4216	-0.3913	4.9111	11.92	
-0.0484	-1.0597	3.5333	13.59	
1.4978	2.9955	5.6177	16.98	
1.9409	3.3233	5.3060	18.58	
2.5161	4.5343	4.5739	14.90	
2.6834	5.1538	3.8993	18.03	
2.6859	5.2414	3.5885	18.51	
<> 13 F(N) Minima :				
-0.5249	1.6155	4.6668	-47.69	
-1.3070	2.2637	4.5275	-45.62	
2.8614	-2.0789	-2.9678	-31.33	
-3.4167	1.0124	2.5040	-26.32	
-0.7780	-4.8993	-0.4346	-30.95	
-1.2131	5.3508	4.2164	-21.65	
-0.6484	-4.9585	-1.1099	-31.95	
0.9079	5.7325	1.0276	-50.35	
0.0536	-4.2250	-2.7041	-41.20	
-0.0608	1.3326	4.4429	-48.57	
3.4583	-1.4318	0.1637	-12.76	
-0.6375	-4.8517	-1.5491	-33.01	
1.0471	-3.6236	-3.4752	-41.55	
<> ParaSurf used 7.26 seconds CPU time				

Figure 14 ParaSurf™ output for trimethoprim, 1, using a spherical-harmonic surface

After printing the program options, ParaSurf'12™ prints the shift in coordinates of the center and the RMSD fits for the surface requested by the **translate** option. For speed, these fits use a lower number of surface points than the full fits that follow and are only calculated up to order six. The translated spherical-harmonic coefficients are printed in the output SDF. ParaSurf'12™ then moves on to fit the calculated shrink-wrap surface at full resolution for each of the local properties. It lists the root-mean-square deviations (RMSDs) for the surface points as a function of the order of the spherical-harmonic expansion, first for the geometry of the surface and then for each of the five local properties. The RMSD values give an idea of how well each order of the spherical-harmonic expansion fits the calculated shrink-wrap surface or the relevant property. The highest order used by ParaSurf™ is 15 for the surface itself and 20 for each property.

The descriptor table is then printed. For molecules with no surface areas with positive EA_{L+} , $\sigma^2_{EA_{L+}}$ is set to zero. The descriptors are those described in **Table 1**.



The spherical-harmonic hybridization coefficients are then listed for the shape and the five local properties. The coefficients are listed by increasing l starting from zero. The standard rotationally invariant fingerprint (RIF) [37] is printed. Note that the individual RIF-values correspond to the square roots of the hybridization coefficients from the tables above and that the RIF definition has been expanded to include hybridization coefficients of the field normal to the surface (the last 13 elements).

The table of atomic surface properties is derived by first finding the atom that contributes most (according to a Coulson analysis) to the electron density for each surface point. The point is then assigned to this atom and the maxima and minima in the MEP, IE_L , EA_L and F_N as well as the mean local polarizability for the points assigned to each atom are calculated. Note that, because of the fitting procedure, the values reported in this table may contain spurious ones if the fitted surface comes particularly close to an atom (or does not approach it). This situation is generally recognizable from the RMSD values printed for the fit. The surface used to calculate the descriptors and atomic-surface properties is the fitted spherical-harmonic surface of order 15.

The maxima and minima of the local properties selected according to the criteria outlined in Ref. [34] are then listed. These points are defined by their Cartesian coordinates and the corresponding values of the local property. In this example, no significant maxima and minima were found for the field normal to the surface. Generally, more maxima and minima are found for isodensity surfaces than for spherical-harmonic ones.

3.5.2 For a marching-cube surface

Figure 15 shows the output for a calculation using the options **surf=cube** for trimethoprim.

```
ParaSurf™ output for trimethoprim using a marching-cube surface.

<> ParaSurf'12, Revision 410
<> Copyright (c) 2006-2012 Friedrich-Alexander-Universitaet Erlangen-Nuernberg
     and Cepos InSilico Ltd. All rights reserved.

<> Input = trimethiprim.sdf

<<>> Molecule      1 of      1 <<>>

<> Program options :

Using marching-cube isodensity surface
Surface fitting turned off
Using an isodensity surface contour
Isodensity value =  0.7000E-02 electrons/Angstrom***3
Triangulation mesh =    0.20 Angstrom
Using multipole electrostatics

<> AM1   calculation for Trimethoprim
<> Number of triangles = 12956
<> Number of unique points : 6483

<> Property ranges:
Density   :  0.6700E-02 to  0.7334E-02
IE(1)     :        312.99 to      539.60
EA(1)     :       -112.06 to     -19.89
MEP       :        -64.18 to      28.35
Alpha(1)  :        0.2319 to     1.5629
```

**ParaSurf™ output for trimethoprim using a marching-cube surface.**

Field(N) :	-96.37 to 62.75
<> Descriptors :	
Dipole moment :	1.2467 Debye
Dipolar density :	0.4108E-02 Debye.Angstrom**-3
Molecular pol. :	128.5408 Angstrom**3
Molecular weight :	290.32
Globularity :	0.6828
Total surface area :	319.86 Angstrom**2
Molecular volume :	303.48 Angstrom**3
Most positive MEP :	28.35 kcal/mol
Most negative MEP :	-64.18 kcal/mol
Mean +ve MEP :	10.17 kcal/mol
Mean -ve MEP :	-19.06 kcal/mol
Mean MEP :	-3.76 kcal/mol
MEP range :	92.53 kcal/mol
MEP +ve Variance :	41.19 (kcal/mol)**2
MEP -ve Variance :	267.51 (kcal/mol)**2
MEP total variance :	308.70 (kcal/mol)**2
MEP balance parameter:	0.1156
MEP balance*variance :	35.6945 kcal/mol
MEP skewness :	-1.0392
MEP kurtosis :	0.6468
Integral MEP :	-1121.97 kcal.Angstrom**2/mol
Maximum IE(1) :	539.60 kcal/mol
Minimum IE(1) :	312.99 kcal/mol
Mean IE(1) :	419.37 kcal/mol
IE(1) range :	226.62 kcal/mol
IE(1) variance :	1693.06 (kcal/mol)**2
IE(1) skewness :	0.5370
IE(1) kurtosis :	-0.2654
Integral IE(1) :	5814.56 eV.Angstrom**2
Maximum EA(1) :	-19.89 kcal/mol
Minimum EA(1) :	-112.06 kcal/mol
Mean +ve EA(1) :	0.00 kcal/mol
Mean -ve EA(1) :	-86.90 kcal/mol
Mean EA(1) :	-86.90 kcal/mol
EA(1) range :	92.17 kcal/mol
EA(1) +ve variance :	0.00 (kcal/mol)**2
EA(1) -ve variance :	363.34 (kcal/mol)**2
EA(1) total variance :	363.34 (kcal/mol)**2
EA(1) skewness :	1.3463
EA(1) kurtosis :	1.2524
Integral EA(1) :	-1215.56 eV.Angstrom**2
EA(1) balance param. :	0.0000
Fraction pos. EA(1) :	0.0000 (= 0.00 Angstrom**2)
Max. local Eneg. :	231.55 kcal/mol
Min. local Eneg. :	108.12 kcal/mol
Mean local Eneg. :	166.23 kcal/mol
Local Eneg. range :	123.43 kcal/mol
Local Eneg. variance :	417.35 (kcal/mol)**2
Local Eneg. skewness :	0.33
Local Eneg. kurtosis :	-0.11
Integral local Eneg. :	2299.50 eV.Angstrom**2
Max. local hardness :	312.13 kcal/mol
Min. local hardness :	191.56 kcal/mol
Mean local hardness :	253.13 kcal/mol
Local hard. range :	120.57 kcal/mol
Local hard. variance :	610.84 (kcal/mol)**2
Local hard. skewness :	0.10
Local hard. kurtosis :	-0.25
Integral local Hard. :	3515.06 eV.Angstrom**2
Maximum alpha(1) :	1.563 Angstrom**3
Minimum alpha(1) :	0.2319 Angstrom**3



ParaSurf™ output for trimethoprim using a marching-cube surface.

```

Mean alpha(1)      : 0.6118    Angstrom**3
Alpha(1) range     : 1.331     Angstrom**3
Variance in alpha(1) : 0.8362E-01 Angstrom**6
Alpha(1) skewness   : 0.7110
Alpha(1) kurtosis   : -0.6335
Integral Alpha(1)   : 192.581    Angstrom**5

Maximum field normal : 62.75 kcal/mol.Angstrom
Minimum field normal : -96.37 kcal/mol.Angstrom
Mean field           : -0.24 kcal/mol.Angstrom
Field range          : 159.12 kcal/mol.Angstrom
Total field variance : 526.01 (kcal/mol.Angstrom)**2
+ve field variance   : 70.61 (kcal/mol.Angstrom)**2
-ve field variance   : 604.96 (kcal/mol.Angstrom)**2
Field balance param. : 0.09
Field skew           : 2.30
Field kurtosis        : 3.507
Integral F(N)         : 14.38      kcal.Angstrom/mol
Integral F(N +ve)     : 2555.      kcal.Angstrom/mol
Integral F(N -ve)     : -2541.     kcal.Angstrom/mol
Integral |F(N)|       : 5096.      kcal.Angstrom/mol

```

<> Atomic surface properties:

Atom	Area	MEP		IE(1)		EA(1)		mean	Field(N)		Eneg(L)		Hard(L)		
		max	min	max	min	max	min		max	min	max	min	max	min	
C	1	3.633	14.36	-27.83	492.02	425.67	-81.81	-98.38	0.729	11.25	-21.64	201.75	164.72	290.27	260.96
O	2	5.872	-16.60	-61.42	517.24	376.50	-65.77	-85.49	1.083	-1.05	-73.60	224.26	154.07	292.99	222.43
C	3	6.547	2.50	-44.56	505.88	359.14	-22.48	-87.04	0.908	9.75	-28.36	213.01	152.78	292.88	192.76
C	4	5.042	11.25	-15.57	482.09	357.89	-27.99	-103.87	0.839	-1.35	-18.32	197.72	149.42	286.32	193.04
C	5	2.637	0.36	-15.56	478.00	380.11	-39.08	-99.66	0.854	0.21	-18.11	196.09	156.43	281.92	209.82
C	6	0.736	2.62	-6.47	484.34	448.62	-90.49	-102.29	0.787	1.54	-11.13	194.67	174.65	289.95	273.97
C	7	3.866	-1.32	-16.72	471.01	352.79	-29.26	-97.15	0.914	8.05	-20.00	190.22	145.74	281.42	197.86
C	8	7.747	8.21	-30.02	485.19	370.16	-27.37	-100.39	0.838	9.32	-27.15	195.38	153.82	289.81	201.38
N	9	8.821	-6.66	-61.59	513.72	315.67	-68.02	-106.03	1.220	-1.63	-93.58	211.59	108.12	302.12	205.46
C	10	9.463	10.11	-34.05	533.20	394.25	-35.99	-99.27	0.895	16.99	-30.32	223.14	168.66	310.05	215.12
N	11	3.906	18.92	-40.19	525.07	373.54	-56.93	-87.16	0.924	12.75	-60.56	227.36	149.70	297.71	223.84
N	12	8.304	-7.20	-58.48	498.61	312.99	-67.59	-105.70	1.114	-3.98	-96.37	204.03	109.56	294.58	199.79
C	13	7.528	8.25	-31.03	528.52	376.72	-29.28	-100.87	0.883	15.91	-29.86	224.59	163.15	306.16	204.32
N	14	3.839	16.28	-39.24	517.43	374.79	-56.71	-89.93	0.939	14.21	-65.25	223.29	150.78	294.14	223.72
C	15	6.243	2.69	-15.27	471.78	355.95	-27.14	-91.73	0.831	3.88	-19.40	190.59	158.13	281.19	192.04
C	16	6.375	-1.57	-42.44	496.09	359.34	-19.89	-90.31	0.891	7.12	-26.86	208.41	158.55	288.34	191.56
O	17	4.315	-12.61	-52.21	517.66	379.79	-66.11	-94.14	0.982	-3.07	-65.91	222.62	148.11	295.03	223.59
C	18	3.756	13.44	-26.67	492.59	427.45	-76.64	-97.98	0.764	10.83	-30.90	205.62	164.89	290.68	261.83
C	19	5.797	-4.74	-53.09	525.33	337.63	-23.77	-96.21	0.962	0.65	-42.42	222.15	142.33	303.17	193.97
O	20	5.835	-37.88	-64.18	524.48	358.58	-72.15	-95.74	1.098	9.43	-73.93	222.32	136.94	302.16	221.62
C	21	4.118	4.81	-44.55	495.26	412.37	-90.33	-111.11	0.769	12.48	-34.16	202.46	151.69	292.79	259.79
H	22	13.548	13.64	-26.76	467.39	383.82	-84.51	-97.89	0.437	17.14	-5.25	189.48	145.14	277.91	238.67
H	23	11.587	18.01	-5.94	480.28	382.02	-81.52	-95.71	0.409	10.75	-7.60	199.38	145.39	281.10	236.63
H	24	11.815	16.82	-11.32	468.75	382.79	-75.15	-95.47	0.409	14.32	-3.06	193.24	145.38	278.42	237.41
H	25	6.767	18.89	0.75	451.49	404.00	-79.12	-101.07	0.393	15.88	-5.11	179.17	153.86	274.47	241.64
H	26	10.613	14.79	-4.33	469.02	375.28	-92.05	-102.60	0.418	11.05	-10.08	186.54	139.27	282.48	236.00
H	27	9.786	13.28	-16.03	476.30	373.94	-90.75	-102.75	0.424	16.09	-10.21	192.20	137.74	284.11	236.20
H	28	11.191	15.84	-24.19	453.71	393.19	-60.41	-100.67	0.431	19.13	-13.24	176.52	150.21	277.19	230.01
H	29	12.557	28.32	-32.30	537.65	465.98	-73.96	-111.21	0.370	58.46	-39.59	231.55	177.41	312.13	278.59
H	30	12.354	28.35	-28.15	538.82	465.58	-77.89	-111.25	0.349	62.75	-29.94	230.46	177.29	311.69	274.57
H	31	12.658	26.33	-29.11	539.60	461.88	-75.50	-112.06	0.364	50.74	-27.22	230.86	174.94	309.67	278.15
H	32	7.359	26.07	-11.11	539.23	414.89	-75.34	-106.39	0.394	32.98	-26.16	230.48	157.36	308.74	257.53
H	33	6.884	15.97	-8.00	446.99	389.30	-77.35	-99.71	0.450	14.37	-13.14	175.53	148.86	271.89	237.01
H	34	13.443	13.65	-24.83	466.77	384.57	-85.30	-96.78	0.424	15.59	-26.06	189.59	145.87	277.17	238.69
H	35	11.745	15.14	-8.63	467.45	382.69	-72.65	-95.71	0.412	13.76	-0.45	189.59	145.33	277.86	237.36
H	36	11.570	14.77	-5.76	474.04	381.97	-77.50	-96.16	0.421	12.80	-1.84	198.27	145.06	278.15	236.91
H	37	13.434	5.48	-36.63	446.92	370.92	-97.15	-109.63	0.420	10.58	-16.68	174.74	133.24	273.45	237.64
H	38	11.343	4.12	-36.90	453.74	365.57	-92.78	-109.05	0.439	12.36	-32.90	175.88	129.49	277.86	236.09
H	39	13.105	4.36	-33.58	487.05	365.53	-85.22	-108.13	0.427	25.62	-10.59	200.91	130.38	286.14	235.15

Total 316.139

<> Stationary points on the molecular surface (A. Jakobi, H. Mauser and T. Clark, J. Mol. Model., 2008, 14, 547-558)

x	y	z	value
<> 13 MEP Maxima :			
4.4569	0.9397	-3.2568	13.65

**ParaSurf™ output for trimethoprim using a marching-cube surface.**

-0.9431	3.5397	-2.9285	15.84
-5.1681	-4.1904	-2.2285	13.64
1.6569	1.8397	-2.5118	15.97
-5.2189	-3.5960	-2.0285	13.79
-4.3431	-1.6853	-1.1618	18.89
3.6569	0.7480	-0.8285	10.06
-3.3931	-0.3603	-0.2618	12.48
-0.7431	-0.4686	0.3715	11.75
-1.9181	-1.4603	1.3548	26.07
-1.9431	-1.4103	1.6933	26.33
1.2069	6.1397	2.4298	21.06
1.2569	4.7204	3.8715	28.35
<>	5 MEP Minima :		
0.4819	-3.8270	-4.1785	-60.72
2.2944	-2.2603	-3.5285	-52.21
-0.4931	-4.2603	-3.2436	-64.18
0.2569	5.3397	0.0840	-61.59
-1.2431	1.7397	3.3715	-58.48
<>	10 IEL Maxima :		
1.6861	-2.6103	-3.4142	525.3
-0.5431	-4.2436	-2.2285	524.5
1.1569	4.5115	1.3715	519.8
-1.8214	-1.4103	1.5715	526.5
-2.7431	-1.2436	1.5715	539.6
-0.6431	5.3397	1.5441	533.2
-2.4431	1.3215	2.5715	533.5
0.4569	2.5397	3.0715	521.6
-1.2931	3.3564	3.4048	533.1
0.6726	5.1397	4.0715	538.8
<>	3 IEL Minima :		
0.0569	5.4397	0.1848	315.7
-1.4431	1.9137	3.3715	313.4
-0.9514	1.6746	3.3715	313.0
<>	6 EAL Maxima :		
-1.4714	-2.0603	-3.9285	-22.48
0.0069	-1.2520	-4.1285	-19.89
-2.5431	-0.8270	-3.5285	-27.99
0.8569	-1.1103	-0.6285	-25.12
-2.4931	3.9397	-0.7067	-27.37
-0.7431	-2.4270	-0.4785	-28.56
<>	16 EAL Minima :		
1.4619	-4.8103	-3.9285	-107.3
1.8319	-4.4936	-3.9285	-107.7
2.1569	-4.2603	-3.8285	-108.6
2.6569	-3.7186	-3.4285	-109.8
0.2569	-5.4603	-2.4285	-111.1
3.4569	-2.7603	-1.3952	-106.6
1.5286	-2.8603	-0.5285	-107.4
-2.5931	-1.5603	0.2298	-105.9
-1.0431	-1.0603	-0.0035	-106.4
0.9569	3.1397	0.2798	-106.0
-2.1431	-1.6436	0.3715	-106.0
0.2569	5.3397	0.8215	-105.6
0.4402	1.7397	1.3465	-105.7
1.2569	5.9397	1.9557	-111.2
-1.7598	-0.3603	3.3715	-112.1
0.2569	3.1397	4.6715	-111.3
<>	3 Alpha(1) Maxima :		
-0.7431	-3.7603	-3.4785	1.339
-0.8648	-3.8603	-3.2785	1.370
0.3370	5.3397	0.4215	1.563
<>	54 Alpha(1) Minima :		
2.0569	0.1397	-4.9285	0.3178
2.2569	0.2447	-4.9785	0.3160
-3.8514	-3.4603	-4.3285	0.3187
-3.3981	-3.2603	-4.3285	0.3178
4.1569	-0.6936	-4.4285	0.3231
2.1402	-5.0603	-3.9285	0.3183

**ParaSurf™ output for trimethoprim using a marching-cube surface.**

4.8069	0.5397	-3.4285	0.3797
-2.3931	1.5347	-3.7285	0.3309
-2.7431	1.7647	-3.6118	0.3267
-1.1431	2.0480	-3.5285	0.3534
1.4319	1.7230	-3.3918	0.3390
-3.0955	-5.4603	-3.3785	0.3368
3.0569	-5.0603	-3.3285	0.3313
-5.1431	-2.7603	-3.2152	0.3362
-5.0431	-2.4103	-2.9952	0.3331
-3.1264	2.1647	-3.2285	0.3250
-2.7431	-5.6436	-2.8535	0.3358
3.4569	-3.3103	-2.6285	0.3429
4.9569	0.1623	-2.5785	0.3595
-0.2598	3.1397	-2.7285	0.3384
-2.7223	-5.7603	-2.2545	0.3337
3.0569	-5.7603	-2.2285	0.3457
-4.1431	-1.0103	-2.5285	0.3418
4.6569	-0.9695	-2.2285	0.3184
0.0902	3.3397	-2.5785	0.3334
-1.8431	4.3454	-2.2285	0.3198
2.0652	-6.2603	-1.8285	0.3643
-4.1431	1.5573	-2.1285	0.3377
2.1069	1.7397	-1.6285	0.3607
-3.9064	-5.5603	-1.2285	0.3498
-0.3431	-4.9603	-1.1118	0.3573
-4.8931	-2.4603	-1.0285	0.3431
1.6569	1.1897	-1.0285	0.3529
-0.3264	-4.2153	-0.6285	0.3554
3.1361	-3.5493	-0.6285	0.3377
3.1319	-0.0603	-0.6285	0.3092
1.3402	-5.6603	-0.4285	0.3542
-3.3931	-0.4603	-0.0335	0.2865
-3.9431	1.5340	-0.3285	0.3554
1.6402	-4.4353	0.1715	0.3239
-3.5931	-3.8603	0.0715	0.3174
-3.2931	-3.6603	0.0715	0.3131
-1.3107	-1.3603	0.5257	0.2593
-0.7431	-0.4686	0.3715	0.3961
1.7569	4.9397	2.0882	0.2451
0.0569	6.0897	1.9382	0.2527
-0.8431	-0.6603	2.4215	0.2322
0.2569	6.2397	2.9715	0.2828
-2.6431	0.3590	3.1715	0.2427
1.6569	4.8680	3.2590	0.2709
1.5569	5.1897	3.3715	0.2611
1.2569	3.2397	3.7998	0.2319
-0.7431	4.0397	4.5715	0.2467
0.4069	4.5397	4.6715	0.2668
 <> 28 F(N) Maxima :			
1.1569	-0.0803	-4.2285	13.38
-2.7431	-2.6803	-3.9285	13.90
4.9569	-0.4404	-3.8035	13.94
-2.2181	2.5397	-3.6285	10.90
-3.0880	-5.7603	-3.0285	13.71
2.4902	-2.8103	-3.1285	10.38
3.9402	-1.7603	-3.2535	14.07
-4.3431	-5.8436	-2.6285	13.91
-3.6848	-5.9603	-2.8785	13.91
-2.6431	-5.6603	-2.6118	14.22
-0.9431	4.5397	-2.6285	18.65
1.2569	2.0897	-1.7902	14.21
-0.4431	-4.3770	-1.6285	22.72
-0.4931	-4.1653	-1.4285	23.01
-4.2931	-1.0603	-1.0285	15.70
2.0569	0.5730	-0.6785	12.24
-4.7431	-2.8445	-0.3285	10.19
-3.6431	0.8897	-0.0285	13.70
-1.6931	-1.5603	0.6498	31.72
0.4569	2.3397	0.6215	11.69
-2.1264	3.5397	1.1715	15.99

**ParaSurf™ output for trimethoprim using a marching-cube surface.**

0.8152	5.2897	1.2715	44.97
1.6069	5.7397	2.3715	34.29
-1.7014	-1.1603	2.7715	31.95
-1.2931	0.5397	3.0715	40.69
-0.3681	2.5564	3.5715	43.03
0.3402	3.7397	4.8715	32.81
0.6819	3.7197	4.7548	33.42
<>	16 F(N) Minima	:	
-0.0157	-4.2603	-3.8285	-72.08
0.7684	-3.6603	-4.2118	-72.28
2.5018	-2.2603	-3.4285	-64.88
-2.4264	-0.6603	-3.5285	-15.74
-0.5931	0.5397	-3.7285	-15.80
3.0402	-2.0853	-2.2285	-58.31
2.8569	-2.0153	-1.8285	-58.39
-1.5256	-4.6103	-1.4285	-69.79
-1.5431	-2.0103	-0.2285	-16.73
-0.0014	-0.0603	-0.2785	-18.93
0.3569	1.5897	-0.1952	-16.03
-3.0931	2.7397	-0.1702	-19.34
0.2569	5.3397	0.0840	-92.35
-3.5431	0.8373	1.2307	-64.12
-0.6931	1.7397	3.3715	-94.22
-1.3431	4.9397	3.1840	-59.11

<> ParaSurf used 6.02 seconds CPU time

Figure 15 ParaSurf™ output for trimethoprim using a marching-cube surface

The table of RMSD values is no longer printed and the range of the electron-density values for the surface points (a test for the quality of the surface) is closer to the target isodensity value (in this case $0.007 \text{ e}^{-\text{\AA}^3}$) than for the fitted surface. The internal precision used by the program is $\pm 2\%$ of the target isodensity value. The values of the descriptors and the atomic-surface properties are more consistent using the marching-cube surface and are recommended for QSPR and surface-integral applications.



3.5.3 For a job with Shannon entropy

Figure 16 and **Figure 17** show the relevant sections of the output for a calculation using the options `surf=cube` for trimethoprim with the extra `shannon` option, which requests internal and external Shannon entropies using the default statistical background file from the `PARASURF_ROOT` directory. The output is identical to that shown in **Figure 15** except that an additional Shannon entropy block is printed after the descriptors, as shown in **Figure 16**:

```
internalexternal
    Maximum Shannon H      :      0.4229      0.3305      bits Angstrom**-2
    Minimum Shannon H     :      0.0464      0.1296      bits Angstrom**-2
    Mean Shannon H       :      0.2126      0.2186      bits Angstrom**-2
    Variance Shannon H   :      0.0073      0.0013      bits Angstrom**-2
    Molecular Shannon H  :      68.76      69.98      bits
```

Figure 16 Shannon entropy section of the ParaSurf™ output for trimethoprim, 1, using a marching-cube isodensity surface.

If the statistical background file is not found or does not have the correct format, only the “internal” Shannon entropy appears in this table. “Internal” Shannon entropy is calculated using the statistical distribution of the local properties on the surface of the molecule itself as reference, whereas “external” Shannon entropy uses pre-calculated background statistics from a database of drug-like ligands. [41]

Note that external statistics files are only provided for the AM1 Hamiltonian.

The Shannon entropy is also analyzed based on the surfaces assigned to the individual atoms to give the table shown in **Figure 17**:

Shannon entropy section of the ParaSurf™ output for trimethoprim, 1, using a marching-cube isodensity surface

Shannon-entropy analysis :

Atom	Area	Internal				Shannon Entropy				External			
		max	min	mean	total	max	min	mean	total	max	min	mean	total
C	1	3.633	0.2135	0.1191	0.1784	0.6481	0.2589	0.1902	0.2189	0.7952			
O	2	5.872	0.1466	0.0545	0.1120	0.6577	0.2334	0.1350	0.1985	1.1656			
C	3	6.547	0.1954	0.0744	0.1383	0.9054	0.2881	0.1481	0.2100	1.3751			
C	4	5.042	0.2322	0.0982	0.1639	0.8263	0.3080	0.1634	0.2373	1.1966			
C	5	2.637	0.2285	0.1216	0.1634	0.4307	0.3065	0.1978	0.2332	0.6148			
C	6	0.736	0.2277	0.1525	0.1854	0.1365	0.2571	0.2264	0.2415	0.1778			
C	7	3.866	0.2273	0.0856	0.1503	0.5810	0.3133	0.1582	0.2203	0.8519			
C	8	7.747	0.2071	0.1233	0.1623	1.2570	0.3041	0.1672	0.2325	1.8010			
N	9	8.821	0.2147	0.0555	0.1166	1.0287	0.2639	0.1375	0.1943	1.7138			
C	10	9.463	0.1988	0.0877	0.1453	1.3751	0.2781	0.1468	0.2087	1.9747			
N	11	3.906	0.1953	0.1013	0.1297	0.5068	0.2528	0.1556	0.2101	0.8209			
N	12	8.304	0.1955	0.0595	0.1193	0.9905	0.2588	0.1461	0.1960	1.6276			
C	13	7.528	0.1900	0.0922	0.1460	1.0993	0.2746	0.1488	0.2095	1.5768			
N	14	3.839	0.1935	0.0971	0.1296	0.4974	0.2418	0.1475	0.2131	0.8181			
C	15	6.243	0.2357	0.1006	0.1504	0.9391	0.3079	0.1690	0.2229	1.3916			
C	16	6.375	0.2046	0.0788	0.1425	0.9084	0.2821	0.1543	0.2052	1.3081			
O	17	4.315	0.2239	0.0690	0.1249	0.5389	0.2435	0.1566	0.2072	0.8941			
C	18	3.756	0.2255	0.1233	0.1845	0.6931	0.2578	0.1917	0.2224	0.8355			
C	19	5.797	0.2178	0.0676	0.1214	0.7038	0.2909	0.1307	0.1970	1.1422			
O	20	5.835	0.1675	0.0587	0.1190	0.6945	0.2434	0.1609	0.2143	1.2503			
C	21	4.118	0.2069	0.1260	0.1528	0.6292	0.2498	0.1553	0.1911	0.7870			
H	22	13.548	0.3922	0.1129	0.2909	3.9414	0.2870	0.1983	0.2393	3.2424			
H	23	11.587	0.3890	0.1496	0.3034	3.5158	0.2795	0.1795	0.2357	2.7315			
H	24	11.815	0.4071	0.1264	0.3175	3.7516	0.3038	0.1826	0.2457	2.9026			
H	25	6.767	0.3387	0.1741	0.2669	1.8066	0.3114	0.1730	0.2048	1.3856			



Shannon entropy section of the ParaSurf™ output for trimethoprim, 1, using a marching-cube isodensity surface											
H	26	10.613	0.3405	0.1561	0.2671	2.8346	0.2673	0.2038	0.2373	2.5183	
H	27	9.786	0.3281	0.1500	0.2615	2.5588	0.2760	0.1891	0.2388	2.3368	
H	28	11.191	0.4103	0.1267	0.3037	3.3982	0.3079	0.1736	0.2416	2.7035	
H	29	12.557	0.2481	0.1108	0.1692	2.1243	0.2587	0.1401	0.1736	2.1795	
H	30	12.354	0.2736	0.1009	0.1649	2.0367	0.2564	0.1392	0.1724	2.1301	
H	31	12.658	0.2601	0.0908	0.1631	2.0650	0.2512	0.1390	0.1741	2.2040	
H	32	7.359	0.2899	0.0994	0.1792	1.3188	0.2651	0.1393	0.1795	1.3214	
H	33	6.884	0.3384	0.1664	0.2591	1.7834	0.3096	0.1834	0.2267	1.5607	
H	34	13.443	0.4137	0.1221	0.3069	4.1249	0.2824	0.1974	0.2386	3.2078	
H	35	11.745	0.4136	0.1311	0.3171	3.7246	0.3046	0.1880	0.2443	2.8692	
H	36	11.570	0.3938	0.1278	0.3115	3.6039	0.3041	0.1868	0.2443	2.8270	
H	37	13.434	0.3376	0.1440	0.2558	3.4362	0.2668	0.1668	0.2349	3.1552	
H	38	11.343	0.3255	0.1204	0.2391	2.7124	0.2671	0.1680	0.2312	2.6222	
H	39	13.105	0.3345	0.1122	0.2579	3.3800	0.2850	0.1639	0.2382	3.1220	

Figure 17 Shannon entropy section of the ParaSurf™ output for trimethoprim, 1, using a marching-cube isodensity surface

3.5.4 For a job with autocorrelation similarity

In order to calculate, for instance, the autocorrelation similarities between captopril and trimethoprim, first calculate the reference compound (in this case captopril) and request that the autocorrelation functions be written to the ParaSurf™ SDF-output file:

```
parasurf captopril surf=cube autocorr
```

Then calculate the autocorrelations for trimethoprim and their similarities to those of captopril:

```
parasurf trimethoprim surf=cube autocorr=captopril_p.sdf
```

This leads to the following additional output from ParaSurf™

```
<> Surface Autocorrelation vectors written to the SD-File
<> Calculating autocorrelation similarities to captopril_p.sdf
<> Lead molecule = Captopril
<> Individual autocorrelation similarities;
    Shape      MEP(tot)    MEP(+-)    MEP(++)    MEP(--)
0.7833      0.6309      0.7873      0.8122      0.5133      0.9720      0.3303      0.9634
<>                                         Total autocorrelation fingerprint similarity = 0.9716
```

Figure 18 Similarity output using autocorrelation functions. The lead molecule is captopril, which is defined in captopril_p.sdf

The “Total autocorrelation fingerprint similarity” refers to the shape, MEP(+), MEP(++) and MEP(--) autocorrelation functions (a total of 448 bins). It is, however dominated by IE(L) and EA(L) because their numerical values are far larger than the other autocorrelation functions.



3.6 ParaSurf™ SDF-output

The SDF output file (a fixed-format file) contains additional blocks with the information generated by ParaSurf™. These are:

<ParaSurf OPTIONS>

The ParaSurf™ OPTIONS block consists of one line giving the options used in the ParaSurf™ calculation. These are:

```
<surface> <fit> <electrostatic model> <isodensity level>      (a4,2x,a4,2x,a5,2x,f8.3)
```

Where the individual variables can be:

<surface>	WRAP	Shrink-wrap surface
	CUBE	Marching-cube surface
<fit>	NONE	No fitting, unsmoothed marching-cube surface
	ISO	Marching-cube surface corrected to $\pm 2\%$ of the preset-isodensity value
	SPHH	Spherical-harmonic surface fit
<electrostatic model>	NAOPC	NAO-PC electrostatics
	MULTI	Multipole electrostatics
<isodensity level>	<i>n.nn</i>	The target isodensity value in $e^{-}\text{\AA}^{-3}$
<solvent probe radius>		The radius of the solvent probe used to calculate the SES or SAS
<triangulation mesh>		The mesh size used to triangulate the Surface

<MOLECULAR_CENTERS>

The molecular centers block appears only for calculations that use spherical harmonic fits. It includes two lines of the form:

```
"Spherical harmonic center = ", 3f12.6  
"Center of gravity           = ", 3f12.6
```

These blocks give the x, y and z coordinates of the center of the molecule used for the spherical-harmonic fit and the center of gravity, respectively. These two centers are usually identical, but may be different if the center of gravity lies outside the molecule (e.g. for U-shaped molecules).

**<SPHERICAL_HARMONIC_.....>**

The spherical harmonic fits are described in **<SPHERICAL_HARMONIC_.....>** blocks. These blocks all have the same format and vary only in the property described. Each block has the form:

Order = nn	("Order = ",i4)
l (c_l^m)m = -1 to 1	(15, 10f8.4/5x,10f8.4/5x,10f8.4/5x,10f8.4) (One set of coefficients each for l = 1 to 15)
RMSDs: l, RMSD¹, RMSD²	("RMSDs:") (i8, 2f12.8) (One line for each l for l= 1 to 15, where RMSD ¹ is the area-weighted RMSD and RMSD ² the simple RMSD)

There are six such blocks, indicated by the tags:

<SPHERICAL_HARMONIC_SURFACE>

The fitted molecular surface (radial distances) in Å

<SPHERICAL_HARMONIC_MEPR>

The MEP values at the spherical-harmonic surface (l= 20) in kcal mol⁻¹

<SPHERICAL_HARMONIC_IE(l)>

The IE_L values at the spherical-harmonic surface (l= 20) in kcal mol⁻¹

<SPHERICAL_HARMONIC_EA(l)>

The EA_L values at the spherical-harmonic surface (l= 20) in kcal mol⁻¹

<SPHERICAL_HARMONIC_ALPHA(l)>

The α_L values at the spherical-harmonic surface (l= 20) in kcal mol⁻¹

<SPHERICAL_HARMONIC_FIELD(N)>

The FN values at the spherical-harmonic surface (l= 20) in kcal mol⁻¹ Å⁻¹

<ParaSurf Descriptors>

The ParaSurf™ descriptors block lists the calculated descriptors in the following groups:

Molecular:	μ, μ _D , α, MW, G, VOL ("Molecular ",5f10.4,2f10.2)
MEP:	V _{max} , V _{min} , \bar{V}_+ , \bar{V}_- , \bar{V} , ΔV, σ_+^2 , σ_-^2 , σ_{Tot}^2 , ν, σ_{tot}^2 V, γ_1^V , γ_2^V , \int_V ("MEP ",7f10.2/10x, f10.2,5f10.4,2x,g12.6)
IE(l):	I _E _L ^{max} , I _E _L ^{min} , \bar{I}_{E_L} , ΔI _E _L , σ_{IE}^2 , γ_1^{IE} , γ_2^{IE} , \int_{IE} ("IE(l) ",5f10.2,2f10.4/12x,g12.6)
EA(l):	E _A _L ^{max} , E _A _L ^{min} , \bar{E}_{A_L} , E _A _L ⁻ , E _A _L , ΔE _A _L , σ_{EA+}^2 , σ_{EA-}^2 , σ_{EA}^2 , V _{EA} , δA _{EA} ⁺ , A _{EA} ⁺ , γ_1^{EA} , γ_2^{EA} , \int_{EA} ("EA(l) ",7f10.2/2f10.2,2f10.4,f10.2,2f10.4/12x,g12.6)
Eneg(l):	χ _L ^{max} , χ _L ^{min} , $\bar{\chi}_L$, Δχ _L , σ_χ^2 , γ_1^χ , γ_2^χ , \int_χ ("Eneg(l) ",5f10.2,2f10.4/12x,g12.6)
Hard(l):	η _L ^{max} , η _L ^{min} , $\bar{\eta}_L$, Δη _L , σ_η^2 , γ_1^η , γ_2^η , \int_η ("Hard(l) ",5f10.2,2f10.4/12x,g12.6)
Alpha(l):	α _L ^{max} , α _L ^{min} , $\bar{\alpha}_L$, Δα _L , σ_α^2 , γ_1^α , γ_2^α , \int_α ("Alpha(l) ",5f10.2,2f10.4/12x,g12.6)
FN	F _N ^{max} , F _N ^{min} , \bar{F}_N , σ_F^2 , σ_{F+}^2 , σ_{F-}^2 , V _F , $\gamma_1^{F_N}$, $\gamma_2^{F_N}$, \int_{F_N} , $\int_{F_N}^+$, $\int_{F_N}^-$, \int_{F_N} ("Field desc",7f10.4/" ",6f10.4)



Jobs that include Shannon entropy give two extra sets of descriptors:

Shannon(i):	$H_{in}^{\max}, H_{in}^{\min}, \overline{H_{in}}, \sigma_{H_{in}}^2, \int_{H_{in}}$	(<code>"Shannon(i) ", 4f10.4, f10.2, f10.4</code>)
Shannon(e):	$H_{ex}^{\max}, H_{ex}^{\min}, \overline{H_{ex}}, \sigma_{H_{ex}}^2, \int_{H_{ex}}$	(<code>"Shannon(e) ", 4f10.4, f10.2, f10.4</code>)

For calculations using a spherical-harmonic fit, the hybridization coefficients are printed to the `.sdf` file as follows (tag line followed by as many lines with the coefficients as necessary):

<SHAPE HYBRIDS>	(15 coefficients, 6f12.6)
<MEP HYBRIDS>	(20 coefficients, 6f12.6)
<IE (L) HYBRIDS>	(20 coefficients, 6f12.2)
<EA (L) HYBRIDS>	(20 coefficients, 6f12.2)
<ALPHA (L) HYBRIDS>	(20 coefficients, 6f12.8)
<FIELD (N) HYBRIDS>	(20 coefficients, 6f12.4)

The hybridization coefficients are listed in order of increasing *I* from zero, exactly as in the output file.

The atomic surface properties are listed in the atomic order according to the following headings (tag line followed by as many lines with the surface properties as necessary):

<ATOMIC SURFACE AREAS>	Areas	(10f8.4)
<ATOMIC SURFACE MEP MAXIMA>	MEP maxima	(10f8.2)
<ATOMIC SURFACE MEP MINIMA>	MEP minima	(10f8.2)
<ATOMIC SURFACE IE (L) MAXIMA>	IE(I) maxima	(10f8.2)
<ATOMIC SURFACE IE (L) MINIMA>	IE(I) minima	(10f8.2)
<ATOMIC SURFACE EA (L) MAXIMA>	EA(I) maxima	(10f8.2)
<ATOMIC SURFACE EA (L) MINIMA>	EA(I) minima	(10f8.2)
<ATOMIC SURFACE MEAN POL>	Mean pol.	(10f8.4)
<ATOMIC SURFACE FIELD (N) MAXIMA>	FN maxima	(10f8.2)
<ATOMIC SURFACE FIELD (N) MINIMA>	FN minima	(10f8.2)

The properties correspond exactly to those printed in the table of surface properties in the output file.



<PROPERTY MAXIMA and MINIMA>

The ParaSurf™ block for the maxima and minima of the local properties is defined as follows for each property:

Header line (maxima)	Number of maxima for the property: N_{max} , property {MEP, IEL, EAL or Alpha(L)}	(I3,a," Maxima")
Nmax maxima lines	x, y, z , property value	(3f12.4,3x,g10.4)
Header line (minima)	Number of minima for the property: N_{min} , property {MEP, IEL, EAL or Alpha(L)}	(I3,a," Minima")
Nmin minima lines	x, y, z , property value	(3f12.4,3x,g10.4)

<STANDARD RIF>

The rotationally invariant fingerprint [37] is printed as a list of 54 floating point numbers (6g12.6). The first 41 are those defined in Ref. [37] and the last 13 are the square roots of the hybridization coefficients for the normal field from $l=0-12$.

3.6.1 Optional blocks in the SDF-output file

A calculation including Shannon entropy gives two extra lines in the descriptors block of the SDF-output file:

The maximum, minimum, mean, variance and total “internal” Shannon entropies.

“Shannon (i)”

(4f10.4,f10.2,f10.4)

The maximum, minimum, mean, variance and total “external” Shannon entropies (if these are calculated).

“Shannon (e)”

(4f10.4,f10.2,f10.4)



Additionally, extra blocks for the atomic Shannon entropy-related variables are added to the SDF-output after the other atomic-property blocks:

<ATOMIC SURFACE MAXIMUM H (internal)>	
Maximum "internal" Shannon entropies	(10f8.4)
<ATOMIC SURFACE MINIMUM H (internal)>	
Minimum "internal" Shannon entropies	(10f8.4)
<ATOMIC SURFACE MEAN H (internal)>	
Mean "internal" Shannon entropies	(10f8.4)
<ATOMIC SURFACE TOTAL H (internal)>	
Total "internal" Shannon entropies	(10f8.4)

If the external Shannon entropy is also calculated, the following blocks are also written:

<ATOMIC SURFACE MAXIMUM H (external)>	
Maximum "external" Shannon entropies	(10f8.4)
<ATOMIC SURFACE MINIMUM H (external)>	
Minimum "external" Shannon entropies	(10f8.4)
<ATOMIC SURFACE MEAN H (external)>	
Mean "external" Shannon entropies	(10f8.4)
<ATOMIC SURFACE TOTAL H (external)>	
Total "external" Shannon entropies	(10f8.4)

For calculations that include surface autocorrelations, these are written in the following blocks:

<SURFACE AUTOCORRELATION PARAMETERS>	
The number of autocorrelation points	("ncorr = ",i6)
The lower end of the autocorrelation range	("rmin = ",f10.6)
The bin size	("dcorr = ",f10.6)



This block then contains a table that gives all the autocorrelations as a table with the following headings:

Table 8 Column headings and definitions for autocorrelation tables.

Column heading	Contents
R	Reference distance (R in Eq. (18))
shape	Shape autocorrelation
MEP(Tot)	Total MEP autocorrelation
MEP(+-)	MEP +/- autocorrelation
MEP(++)	MEP ++/ autocorrelation
MEP(--)	MEP -/- autocorrelation
IE(L)	IEL autocorrelation
EA(L)	EAL autocorrelation
Alpha(L)	Alpha(L) autocorrelation

The format of the columns is (f8.2, 2x, 8g15.6)

Calculations with spherical-harmonic fits that use the **TRANSLATE** or **TRANSLATE2** options, an additional block with the header

<TRANSLATED SPHERICAL HARMONIC FITS>

is printed. This block consists of nine sets of results (the original center plus eight translated ones) for **TRANSLATE** and 16 for **TRANSLATE2**. The original center is denoted by the header

```
Origin <shiftx> <shifty> <shiftz> <RMSD>
("Origin :",3f12.4,f12.6)'
```

followed by the fitted coefficients (7f12.6). The shifted points are defined in the same way, but are denoted "Point N"

```
("Point ",i2,:",3f12.4,f12.6)
```



3.7 The surface (.psf) file

The **.psf** file can be used to derive properties and descriptors from the ParaSurf™ results. It includes the coordinates and properties of the atoms, surface points and surface triangles in the following format. This format has been extended compared to that used by ParaSurf'11™.

Number of atoms	(i6)
------------------------	------

One line per atom with the atomic surface properties:

Atomic number, x-coordinate, y-coordinate, z-coordinate, atomic surface area, V_{max}, V_{min}, IE_L^{min}, EA_L^{max}, mean polarizability	(i2,3f10.5,f8.3,4f8.2,f8.3)
--	-----------------------------

Number of surface points, total number (N_{models}) of surface-integral models (normal and binned)	(i6,1x,i5)
---	------------

The three-letter codes for the individual models	N _{models} * (1x,a3)
---	-------------------------------

One line per point with the local properties:

x-coordinate, y-coordinate, z-coordinate, MEP, IE_L, EA_L, α_L, atom_L, local value of each model	(3f10.5,3f8.2,f8.4,i6,N _{models} * (2x,g12.4))
--	--

(where atom_L is the atom to which the surface point is assigned)

Number of surface triangles	(i6)
------------------------------------	------

One line per triangle with the ID of the triangle and the local properties:

point #1, point #2, point #3, area, atom_{tri}, normal field	(3i6,f10.5,i6,g12.4)
---	----------------------

(where point #1, 2 and 3 are the numbers of the surface points that make up the triangle and atom_{tri} is the atom to which the triangle is assigned)



3.8 Anonymous SD (.asd) files

The **.asd** file contains only those blocks from the ParaSurf™ output SD file that do not pertain directly to the 2D-molecular structure. Its purpose is to allow a full descriptions of the intermolecular bonding properties of the molecule without revealing its structure. The **.asd** file can only be written from a ParaSurf™ calculation using spherical-harmonic fitting. Its form is:

The SD header line	(A molecular ID number etc.)
The program identifier line	(The normal second line of the SD-file)

And the blocks defined by the following tags:

```
<SPHERICAL_HARMONIC_SURFACE>
<SPHERICAL_HARMONIC_MEPE>
<SPHERICAL_HARMONIC_IE(1)>
<SPHERICAL_HARMONIC_EA(1)>
<SPHERICAL_HARMONIC_FIELD(N)>
<SPHERICAL_HARMONIC_ALPHA(1)>
<SHAPE HYBRIDS>
<MEP HYBRIDS>
<IE(L) HYBRIDS>
<EA(L) HYBRIDS>
<FIELD(N) HYBRIDS>
<ALPHA(L) HYBRIDS>
<STANDARD RIF>
```

<ParaSurf Descriptors>

(The molecular weight and the atomic surface properties are not included because they would allow the molecular formula to be reconstructed. The atoms assigned to each surface point or triangle are also not given.) The format of the descriptors is:

Molecular	$\mu, \mu_D, \alpha, MW, G, A, VOL$ ("Molecular ", 5f10.4, 2f10.2)
MEP	$V_{max}, V_{min}, \bar{V}_+, \bar{V}_-, \bar{V}, \Delta V, \sigma_+^2, \sigma_-^2, \sigma_{tot}^2, v, \sigma_{tot}^2 v, \gamma_1^V, \gamma_2^V, \int_V$ ("MEP ", 7f10.2/10x, f10.2, 5f10.4, 2x, g12.6)
IE(I)	$IE_L^{max}, IE_L^{min}, \overline{IE}_L, \Delta IE_L, \sigma_{IE}^2, \gamma_1^{IE}, \gamma_2^{IE}, \int_{IE}$ ("IE(1) ", 5f10.2, 2f10.4/12x, g12.6)
EA(I)	$EA_L^{max}, EA_L^{min}, \overline{EA}_{L+}, \overline{EA}_{L-}, EA_L, \Delta EA_L, \sigma_{EA+}^2, \sigma_{EA-}^2, \sigma_{EA}^2, \nu_{EA}, \delta A_{EA}^+, A_{EA}^+, \gamma_1^{EA}, \gamma_2^{EA}, \int_{EA}$ ("EA(1) ", 7f10.2/2f10.2, 2f10.4, f10.2, 2f10.4/12x, g12.6)



Eneg(l)	$\chi_L^{\max}, \chi_L^{\min}, \overline{\chi_L}, \Delta\chi_L, \sigma_{\chi}^2, \gamma_1^{\chi}, \gamma_2^{\chi}, \int_{\chi}$ ("Eneg(1) ", 5f10.2, 2f10.4/12x, g12.6)
Hard(l)	$\eta_L^{\max}, \eta_L^{\min}, \overline{\eta_L}, \Delta\eta_L, \sigma_{\eta}^2, \gamma_1^{\eta}, \gamma_2^{\eta}, \int_{\eta}$ ("Hard(1) ", 5f10.2, 2f10.4/12x, g12.6)
Alpha(l)	$\alpha_L^{\max}, \alpha_L^{\min}, \overline{\alpha_L}, \Delta\alpha_L, \sigma_{\alpha}^2, \gamma_1^{\alpha}, \gamma_2^{\alpha}, \int_{\alpha}$ ("Alpha(1) ", 5f10.2, 2f10.4/12x, g12.6)
F_N	$F_N^{\max}, F_N^{\min}, \Delta F_N, \overline{F_N}, \sigma_F^2, \sigma_{F+}^2, \sigma_{F-}^2, \nu_F, \gamma_1^{F_N}, \gamma_2^{F_N}, \int_{F_N}, \int_{F_N}^+, \int_{F_N}^-$ ("Field desc", 7f10.4/" ", 6f10.4)

Jobs that include Shannon entropy give two extra sets of descriptors:

Shannon(i)	$H_{in}^{\max}, H_{in}^{\min}, \overline{H_{in}}, \sigma_{H_{in}}^2, \int_{H_{in}}$ ("Shannon(i) ", 4f10.4, f10.2, f10.4)
Shannon(e)	$H_{ex}^{\max}, H_{ex}^{\min}, \overline{H_{ex}}, \sigma_{H_{ex}}^2, \int_{H_{ex}}$ ("Shannon(e) ", 4f10.4, f10.2, f10.4)

3.8.1 Optional blocks

For calculations that include surface autocorrelations, these are written in the following blocks:

<SURFACE AUTOCORRELATION PARAMETERS>

The number of autocorrelation points	("ncorr = ", i6)
The lower end of the autocorrelation range	("rmin = ", f10.6)
The bin size	("dcorr = ", f10.6)

This block then contains a table that gives all the autocorrelations as a table with the following headings:

Table 9 Column headings and definitions for the autocorrelation table in the output SDF file.

Column heading	Contents
R	Reference distance (R in equation (18))
shape	Shape autocorrelation
MEP(Tot)	Total MEP autocorrelation
MEP(+-)	MEP +/- autocorrelation
MEP(++)	MEP ++/+ autocorrelation
MEP(--)	MEP -/- autocorrelation
IE(L)	IEL autocorrelation
EA(L)	EAL autocorrelation
Alpha(L)	Alpha(L) autocorrelation

The format of the columns is (f8.2, 2x, 8g15.6)



3.9 Grid calculations with ParaSurf™

3.9.1 User-specified Grid

The command

```
parasurf <filename> estat=multi grid=grid.dat
```

instructs ParaSurf™ to read a set of Cartesian coordinates from the file grid.dat and to calculate the \log_{10} of the electron density, the four local properties, the electric field and the derivatives of the local properties ($\log(\rho)$, MEP (=V), IE(I), EA(I) Pol(I), Eneg(I), Hard(I), dV/dx , dV/dy , dV/dz , $d\rho/dx$, $d\rho/dy$, $d\rho/dz$, $d\log(\rho)/dx$, $d\log(\rho)/dy$, $d\log(\rho)/dz$, $dIE(I)/dx$, $dIE(I)/dy$, $dEA(I)/dx$, $dEA(I)/dy$, $dEneg/I/dx$, $dEneg/I/dy$, $dHard/I/dx$, $dHard/I/dy$, $dHard/I/dz$). The format of the file grid.dat (which must be in the same directory as the input) is one line per point containing the x, y and z coordinates in free format, comma-separated, maximum line length 80 with no trailing comma. For instance, the following grid file:

-8.01100 , -13.72910 , -7.91090
-8.01100 , -13.72910 , -6.91090
-8.01100 , -13.72910 , -5.91090
-8.01100 , -13.72910 , -4.91090
-8.01100 , -13.72910 , -3.91090
-8.01100 , -13.72910 , -2.91090
-8.01100 , -13.72910 , -1.91090
-8.01100 , -13.72910 , -0.91090
-8.01100 , -13.72910 , 0.08910
-8.01100 , -13.72910 , 1.08910

Figure 19 Sample grid file

gives the output shown in Figure 20.

Sample grid output file (original)

```
<> Parasurf'12, Revision 410
<> Copyright (c) 2006-2012 Friedrich-Alexander-Universitaet Erlangen-Nuernberg
and Cepos InSilico Ltd. All rights reserved.

<> Input = trimethiprim.sdf

<>>> Molecule      1 of      1 <<>>

<> Program options :
Calculating local properties using grid file grid.txt
Using multipole electrostatics
```

Sample grid output file (.txt)

```
<> ParaSurf'12, Revision 410
<> Copyright (c) 2006-2012 Friedrich-Alexander-Universitaet Erlangen-Nuernberg
and Cepos InSilico Ltd. All rights reserved.
```

```
<> Input = trimethiprim.sdf
```

```
<>>> Molecule      1 of      1 <<>>
```

```
<> Program options :
```

```
Calculating local properties using grid file grid.txt
Using multipole electrostatics
```



```
<> AM1      calculation for Trimethoprim

          x           y           z           log(rho)           MEP       IE(1)       EA(1)       Pol(1)
Eneg(1)   Hard(1)   dv/dx   dv/dy   dv/dz   dRho/dx   dRho/dy   dRho/dz
dlogR/dx   dlogR/dy   dlogR/dz   dIEL/dx   dIEL/dy   dIEL/dz   dEA1/dx   dEA1/dy
dEA1/dz   dEneq/dx   dEneq/dy   dEneq/dz   dHard/dx   dHard/dy   dHard/dz

-8.01100 -13.72910 -7.91090 -21.981311 -0.14 397.68 -93.93 0.4314
151.87    245.80   6.165E-02 -3.937E-04 -3.356E-02 1.787E-22 3.681E-22 2.289E-22
7.432E-01 1.531E+00 9.523E-01 -1.928E+00 2.328E+00 -2.815E+00 4.315E-02 -5.509E-02
6.822E-02 -9.423E-01 1.137E+00 -1.374E+00 -9.855E-01 1.192E+00 -1.442E+00

-8.01100 -13.72910 -6.91090 -21.096656 -0.10 395.21 -93.87 0.4454
150.67    244.54   6.241E-02 -9.348E-03 -4.102E-02 1.426E-21 2.938E-21 1.501E-21
7.739E-01 1.594E+00 8.142E-01 -1.870E+00 1.701E+00 -2.117E+00 4.256E-02 -4.129E-02
5.351E-02 -9.135E-01 8.301E-01 -1.032E+00 -9.561E-01 8.714E-01 -1.085E+00

-8.01100 -13.72910 -5.91090 -20.358433 -0.05 393.44 -93.82 0.4572
149.81    243.63   6.127E-02 -2.024E-02 -4.705E-02 8.086E-21 1.666E-20 6.653E-21
8.016E-01 1.651E+00 6.595E-01 -1.826E+00 1.224E+00 -1.434E+00 4.188E-02 -3.019E-02
3.833E-02 -8.922E-01 5.971E-01 -6.979E-01 -9.341E-01 6.273E-01 -7.362E-01

-8.01100 -13.72910 -4.91090 -19.782872 -0.01 392.34 -93.79 0.4658
149.27    243.07   5.815E-02 -3.235E-02 -5.068E-02 3.130E-20 6.447E-20 1.857E-20
8.245E-01 1.698E+00 4.892E-01 -1.809E+00 8.945E-01 -7.602E-01 4.140E-02 -2.211E-02
2.297E-02 -8.840E-01 4.362E-01 -3.686E-01 -9.254E-01 4.583E-01 -3.916E-01

-8.01100 -13.72910 -3.91090 -19.384276 0.05 391.92 -93.78 0.4701
149.07    242.85   5.324E-02 -4.454E-02 -5.122E-02 7.995E-20 1.647E-19 2.910E-20
8.412E-01 1.733E+00 3.061E-01 -1.830E+00 7.083E-01 -6.999E-02 4.138E-02 -1.717E-02
7.305E-03 -8.942E-01 3.456E-01 -3.134E-02 -9.356E-01 3.628E-01 -3.865E-02

-8.01100 -13.72910 -2.91090 -19.173520 0.10 392.22 -93.78 0.4694
149.22    243.00   4.695E-02 -5.549E-02 -4.847E-02 1.313E-19 2.705E-19 1.766E-20
8.503E-01 1.751E+00 1.143E-01 -1.896E+00 6.709E-01 6.788E-01 4.197E-02 -1.549E-02
-9.132E-03 -9.269E-01 3.277E-01 3.348E-01 -9.688E-01 3.432E-01 3.439E-01

-8.01100 -13.72910 -1.91090 -19.156717 0.14 393.32 -93.80 0.4634
149.76    243.56   3.983E-02 -6.402E-02 -4.281E-02 1.366E-19 2.814E-19 -1.297E-20
8.511E-01 1.753E+00 -8.082E-02 -2.012E+00 8.029E-01 1.542E+00 4.320E-02 -1.731E-02
-2.699E-02 -9.846E-01 3.928E-01 7.576E-01 -1.028E+00 4.101E-01 7.846E-01

-8.01100 -13.72910 -0.91090 -19.334354 0.18 395.36 -93.83 0.4522
150.76    244.60   3.246E-02 -6.935E-02 -3.510E-02 8.993E-20 1.852E-19 -2.917E-20
8.434E-01 1.737E+00 -2.736E-01 -2.180E+00 1.147E+00 2.583E+00 4.486E-02 -2.306E-02
-4.672E-02 -1.068E+00 5.618E-01 1.268E+00 -1.112E+00 5.849E-01 1.315E+00

-8.01100 -13.72910 0.08910 -19.701160 0.21 398.56 -93.89 0.4368
152.33    246.23   2.530E-02 -7.131E-02 -2.641E-02 3.793E-20 7.814E-20 -2.100E-20
8.279E-01 1.705E+00 -4.583E-01 -2.390E+00 1.765E+00 3.857E+00 4.646E-02 -3.306E-02
-6.811E-02 -1.172E+00 8.659E-01 1.895E+00 -1.218E+00 8.989E-01 1.963E+00

-8.01100 -13.72910 1.08910 -20.246709 0.23 403.16 -93.97 0.4182
154.60    248.57   1.868E-02 -7.022E-02 -1.770E-02 1.051E-20 2.166E-20 -8.224E-21
8.058E-01 1.660E+00 -6.304E-01 -2.622E+00 2.727E+00 5.387E+00 4.723E-02 -4.701E-02
-8.951E-02 -1.287E+00 1.340E+00 2.649E+00 -1.335E+00 1.387E+00 2.738E+00

<> ParaSurf used          0.15 seconds CPU time
```

Figure 20 Sample grid output file

The name and the extension (if any) of the grid file are free. Only the output file is written. The units of the local properties are those used in the normal output (*i.e.* V, IEL, and EAL in kcal mol⁻¹, α L in Ångstrom³.



3.9.2 Automatic grids

ParaSurf™ can generate grids automatically for lead compounds in CoMFA®-like procedures. The **grid=auto** option generates a grid around the molecule (with a 0.5 Å margin around the positions of the atoms in each direction) and includes all points for which the electron density is lower than 10⁻² (i.e. for points outside the molecule). The spacing of the grid is set to a default value of 1.0 Å, but can be set to any value up to a maximum of 2.0 Å by the command-line argument **lattice=n.n**, which sets the lattice spacing to **n.n** Å. The grid thus generated is output (with the values of the local properties analogously to a calculation that uses a predefined grid and can be used for other molecules that have been aligned with the lead. An additional output file named **<filename>.p.grid**. There are two further variations of the automatic grid-generation procedure: **grid=auto** excludes any points that are within 0.5 Å of a nucleus, whereas **grid=vdw** excludes all grid points within the van der Waals volume of the molecule and **grid=box** calculates all points regardless of their proximity to a nucleus.



3.10 The SIM file format

SIM files must reside in the ParaSurf™ executable directory and are strictly fixed format. SIM files must be called **<filename>.sim**, where **<filename>** must have exactly three characters. A sample SIM file for a single model (the free energy of solvation in octanol) is shown in Figure 21:

```
><OPTIONS>
surf=cube
fit=isod
estat=multi
iso=0.05
><MODELS>
    1      3
><DGO>
    31.61058
DeltaG(n-Octanol)
kcal/mol
-0.01107    F  1.0      0.0      0.0      1.0      0.0      1.0
  1.6793d-9  F  1.0      0.0      3.0      0.0      0.0      1.0
-2.0407d-10 T  1.0      0.0      1.0      0.0      1.0      1.5
```

Figure 21 Sample surface-integral model (SIM) file

The first line, the OPTIONS tag, is compulsory and takes the form:

<OPTIONS>

The second to fifth lines, also compulsory in the order shown above, give the ParaSurf™ options to be used for the surface-integral model. These options are given in lower case and override conflicting command-line options.

Line 6 must be the MODELS tag with the format

<MODELS>

Line 7 contains the two integers (**Nmodels** and **Maxterms**) that define the number of models given in the file and the maximum number of terms for any one model. The format is:

Nmodels Maxterms

(214)



The remainder of the SIM file consists of **Nmodels** blocks, each of which defines a single model and has the following format:

Model identifier tag

<MOD>

where MOD is a three-letter unique identifier for the model.

Nterms (the number of terms in the model), constant (the constant in the regression equation)	(i4,g12.6)
Model name (for output, maximum 20 characters)	(a20)
Units of the property <i>P</i> (for output, maximum 20 characters)	(a20)
Nterms lines, one per term, giving the definition of the model: Coeff Abs m n o p q r	(d12.6,13,6f8.4)

where each term is defined as:

$$\left[MEP^m \cdot IE_L^n \cdot EA_L^o \cdot \alpha_L^p \cdot \eta_L^q \right]^r \text{ if } \text{Abs} \text{ is false and } \left[|MEP^m \cdot IE_L^n \cdot EA_L^o \cdot \alpha_L^p \cdot \eta_L^q| \right]^r \text{ if } \text{Abs} \text{ is true.}$$

SIM files are only intended to be created by expert users.



3.11 Output tables

The command-line argument `table=<filename>` requests that the 41 descriptors written in the `<ParaSurf DESCRIPTORS>` block of the ParaSurf™ SD-file output are written, one line per molecule, in the file `<filename>`. If `<filename>` already exists, the line for the new molecules will be appended, otherwise a new file will be created and a header line including designations of the descriptors will be written as the first line. All lines in the table file are comma-separated with all blanks (including those in the Molecule ID) removed. The Descriptors in order are:

Table 10 Definitions and order of the descriptors printed to the descriptor table if requested.

Column Header	Symbol ^a	Descriptor
MolID		Molecular ID taken from the first line of the entry for each molecule with all blanks eliminated.
dipole	μ	Dipole moment
dipden	μ_D	Dipolar density
polarizability	α	Molecular electronic polarizability
MWt	MW	Molecular weight
globularity	G	Globularity
totalarea	A	Molecular surface area
volume	VOL	Molecular volume
MEPmax	V_{\max}	Maximum (most positive) MEP
MEPmin	V_{\min}	Minimum (most negative) MEP
meanMEP+	\bar{V}_+	Mean of the positive MEP values
meanMEP-	\bar{V}_-	Mean of the negative MEP values
meanMEP	\bar{V}	Mean of all MEP values
MEPrange	ΔV	MEP-range
MEPvar+	σ_+^2	Total variance in the positive MEP values
MEPvar-	σ_-^2	Total variance in the negative MEP values
MEPvartot	σ_{tot}^2	Total variance in the MEP
MEPbalance	v	MEP balance parameter
var*balance	$\sigma_{tot}^2 v$	Product of the total variance in the MEP and the balance parameter
MEPskew	γ_1^V	Skewness of the distribution of the MEP
MEPkurt	γ_2^V	Kurtosis of the distribution of the MEP
MEPint	\int_V	Integral of the MEP*area over the surface
IELmax	IE_L^{\max}	Maximum value of the local ionization energy
IELmin	IE_L^{\min}	Minimum value of the local ionization energy



Column Header	Symbol ^a	Descriptor
IELbar	\overline{IE}_L	Mean value of the local ionization energy
IELrange	ΔIE_L	Range of the local ionization energy
IELvar	σ_{IE}^2	Variance in the local ionization energy
IELskew	γ_1^{IE}	Skewness of the distribution of IE(L)
IELkurt	γ_2^{IE}	Kurtosis of the distribution of IE(L)
IELint	\int_{IE}	Integral of the IE(L)*area over the surface
EALmax	EA_L^{\max}	Maximum of the local electron affinity
EALmin	EA_L^{\min}	Minimum of the local electron affinity
EALbar+	\overline{EA}_{L+}	Mean of the positive values of the local electron affinity
EALbar-	\overline{EA}_{L-}	Mean of the negative values of the local electron affinity
EALbar	\overline{EA}_L	Mean value of the local electron affinity
EALrange	ΔEA_L	Range of the local electron affinity
EALvar+	σ_{EA+}^2	Variance in the local electron affinity for all positive values
EALvar-	σ_{EA-}^2	Variance in the local electron affinity for all negative values
EALvartot	$\sigma_{EA_{tot}}^2$	Sum of the positive and negative variances in the local electron affinity
EALbalance	ν_{EA}	Local electron affinity balance parameter
EALfraction+	δA_{EA}^+	Fraction of the surface area with positive local electron affinity
EALarea+	A_{EA}^+	Surface area with positive local electron affinity
EALskew	γ_1^{EA}	Skewness of the distribution of the MEP
EALKurt	γ_2^{EA}	Kurtosis of the distribution of the MEP
EALint	\int_{EA}	Integral of the MEP*area over the surface
POLmax	α_L^{\max}	Maximum value of the local polarizability
POLmin	α_L^{\min}	Minimum value of the local polarizability
POLbar	$\overline{\alpha}_L$	Mean value of the local polarizability
POLrange	$\Delta \alpha_L$	Range of the local polarizability
POLvar	σ_{α}^2	Variance in the local polarizability
POLskew	γ_1^{α}	Skewness of the distribution of the local polarizability
POLkurt	γ_2^{α}	Kurtosis of the distribution of the local polarizability
POLint	\int_{α}	Integral of the $\alpha(L)^*$ area over the surface
ENEGmax	χ_L^{\max}	Maximum of the local electronegativity
ENEGmin	χ_L^{\min}	Minimum of the local electronegativity
ENEGbar	$\overline{\chi}_L$	Mean value of the local electronegativity



Column Header	Symbol ^a	Descriptor
ENEGrange	$\Delta\chi_L$	Range of the local electronegativity
NEGvar	σ_χ^2	Variance in the local electronegativity
NEGskew	γ_1^χ	Skewness of the distribution of the local electronegativity
NEGkurt	γ_2^χ	Kurtosis of the distribution of the local electronegativity
NEGint	\int_χ	Integral of the $\chi(L)^*$ area over the surface
HARDmax	η_L^{\max}	Maximum of the local electronegativity
HARDmin	η_L^{\min}	Minimum of the local electronegativity
HARDbar	$\overline{\eta}_L$	Mean value of the local electronegativity
HARDrange	$\Delta\eta_L$	Range of the local electronegativity
HARDvar	σ_η^2	Variance in the local electronegativity
HARDskew	γ_1^η	Skewness of the distribution of the local electronegativity
HARDkurt	γ_2^η	Kurtosis of the distribution of the local electronegativity
HARDint	\int_η	Integral of the $\chi(L)^*$ area over the surface
FNmax	F_N^{\max}	Maximum value of the field normal to the surface
FNmin	F_N^{\min}	Minimum value of the field normal to the surface
FNrange	ΔF_N	Range of the field normal to the surface
FNmean	\overline{F}_N	Mean value of the field normal to the surface
FNvartot	σ_F^2	Variance in field normal to the surface
FNvar+	σ_{F+}^2	Variance in the field normal to the surface for all positive values
FNvar-	σ_{F-}^2	Variance in the field normal to the surface for all negative values
FNbal	ν_F	Normal field balance parameter
FNskew	$\gamma_1^{F_N}$	Skewness of the field normal to the surface
FNkurt	$\gamma_2^{F_N}$	Kurtosis of the field normal to the surface
FNint	\int_{F_N}	Integrated field normal to the surface over the surface
FN+	$\int_{F_N}^+$	Integrated field normal to the surface over the surface for all positive values
FN-	$\int_{F_N}^-$	Integrated field normal to the surface over the surface for all negative values
FNabs	$\int_{ F_N }$	Integrated absolute field normal to the surface over the surface

^aSymbols as used in **Section 1.9**



If the Shannon entropy is calculated, the following additional descriptors are added:

Table 11 Additional descriptors (Shannon entropy)

Column Header	Symbol	Descriptor
SHANImax	H_{in}^{\max}	Maximum internal Shannon entropy
SHANImin	H_{in}^{\min}	Minimum internal Shannon entropy
SHANIvar	$\sigma_{H(in)}^2$	Variance of the internal Shannon entropy
SHANIbar	\bar{H}_{in}	Mean internal Shannon entropy
SHANItot	H_{in}^{tot}	Total internal Shannon entropy

and if the external Shannon entropy is also calculated

SHANE \max	H_{ex}^{\max}	Maximum external Shannon entropy
SHANE \min	H_{ex}^{\min}	Minimum external Shannon entropy
SHANEvar	$\sigma_{H(ex)}^2$	Variance of the external Shannon entropy
SHANEbar	\bar{H}_{ex}	Mean external Shannon entropy
SHANEtot	H_{ex}^{tot}	Total external Shannon entropy
SHANEtot	H_{ex}^{tot}	Total external Shannon entropy



3.12 Binned SIM descriptor tables

If the option `desfile=<filename>` is used, a user-defined file with binned SIM-descriptors is written. The bin limits are taken from installed models using the command-line option `desmodel=<code>`, where `<code>` is the model code taken from **Table 3**. If the table file does not exist, a new one with a header line will be written, otherwise the results for the current molecule will be appended. The descriptors are denoted by a two- or three-letter code to denote the property followed by the number of the descriptor (currently 1-12). The letter codes are:

Table 12 Letter codes (Binned SIM descriptor tables)

Code	Property	Code	Property	Code	Property
MEP	MEP	IEL	IE _L	EAL	EA _L
POL	α _L	FN	F _N	HD	η _L
ENG	χ _L	MI	MEP×IE _L	MA	MEP×EA _L
MP	MEP×α _L	MF	MEP×F _N	MH	MEP×η _L
ME	MEP×χ _L	IA	IE _L ×EA _L	IP	IE _L ×α _L
IF	IE _L ×F _N	IH	IE _L ×η _L	IE	IE _L ×χ _L
AP	EA _L ×α _L	AF	EA _L ×F _N	AH	EA _L ×η _L
AE	EA _L ×χ _L	PF	α _L ×F _N	PH	α _L ×η _L
PE	α _L ×χ _L	FH	F _N ×η _L	FE	F _N ×χ _L
HE	η _L ×χ _L				

The descriptor file is comma-separated.



3.13 Autocorrelation fingerprint and similarity tables

If the option `aclist=<filename>` is used, a user-defined file with the autocorrelation fingerprint is written. If this file does not exist, it is created and the header line written, otherwise entries are appended. The ASCII file is comma-separated and contains the molecular identifier followed by 448 binned autocorrelation values in the order Shape, MEP(+), MEP(++) , MEP(--), IE(L), EA(L), Alpha(L) (64 bins each).

The option `aclist=<filename>` requests a user-defined file with the autocorrelation similarities to the lead compound defined using the `autocorr=<filename>` keyword. If this file does not exist, it is created and the header line written, otherwise entries are appended. The ASCII file has a fixed format. The header line is

```
Molid  Shape  MEP(tot)  MEP(+-)  MEP(++)  MEP(--)  IE(1)  EA(1)  Alpha(1)  Fingerprint
```

and the similarities are written in format (a20,9f10.4). If the molecular identifier is longer than 20 characters, it will be truncated. The "Fingerprint" similarity considers all 448 autocorrelation values (but see [Section 1.15](#)).

3.14 Shared files

The `Vhamil.par` and `SIM` files are accessed in shared, read-only mode so that multiple ParaSurf™ jobs can access the same files.



4 TIPS FOR USING PARASURF'12™

4.1 Choice of surface

ParaSurf™ was originally written to use isodensity surfaces. However, calculations that use a solvent-excluded surface are very much faster than their equivalents with isodensity surfaces and will usually give comparable results. Surface-integral models may benefit from using a solvent-excluded surface with a solvent radius of 0.5-1.0 Å as this appears to be the most relevant surface for many physical properties. Surfaces fitted to spherical-harmonic expansions require more CPU-time than marching-cube surfaces but are essential for fast numerical applications. Again, solvent-excluded shrink-wrap surfaces are faster to calculate than their isodensity equivalents.

4.2 Local properties

The improved local properties implemented in ParaSurf'12™ generally give better QSAR and QSPR models than the earlier ones available up to ParaSurf'11™. It is therefore recommended that new projects use the ParaSurf'12™ local properties.

4.3 QSAR using grids

As outlined in [Section 3.9.2](#), ParaSurf™ can generate a grid for the lead molecule automatically that can then be used for a set of aligned molecules for grid-based QSAR. This procedure has proven to be especially effective for test datasets, especially if the molecules are aligned to a common scaffold, as outlined in [Section 1.1](#).

The automatic grid generated for a lead molecule is now written to the file `<filename>_p.grid` for use with the remainder of the dataset.



5 SUPPORT

5.1 Contact

Questions regarding ParaSurf™ should be sent directly to:

support@ceposinsilico.com

5.2 Error reporting

Some of the routines in ParaSurf™ may detect error conditions that have not yet been encountered in our tests. In this case, an error message will be printed requesting that the input and output files be sent to the programming team at the above e-mail address. We realize that this will not always be possible for confidentiality reasons, but if the details can be sent, we will be able to treat the exception and improve the program.

5.3 Cepos InSilico GmbH

Computer-Chemie-Centrum (CCC)
Nägelsbachstr. 25
91052 Erlangen
Germany

support@ceposinsilico.com
Tel. +49-9131-9704910
Fax. +49-9131-9704911

www.ceposinsilico.com/contact



6 LIST OF TABLES

Table 1	The descriptors calculated by ParaSurf™	16
Table 2	The 28 local properties and products thereof used to construct binned area descriptors.	22
Table 3	Local hydrophobicity models and their model codes (all models use the single CORINA-derived conformations and are trained with the "full" dataset. [30])	22
Table 4	ParaSurf™ command-line options	30
Table 5	ParaSurf™ input and output files	40
Table 6	Hamiltonians and the available electrostatic and polarizability models.	42
Table 7	Content of the EMPIRE™ binary wavefunction file (.vwf)	44
Table 8	Column headings and definitions for autocorrelation tables.	66
Table 9	Column headings and definitions for the autocorrelation table in the output SDF file.	69
Table 10	Definitions and order of the descriptors printed to the descriptor table if requested.	75
Table 11	Additional descriptors (Shannon entropy)	78
Table 12	Letter codes (Binned SIM descriptor tables)	79



7 LIST OF FIGURES

Figure 1	Marching-cube (left) and shrink-wrap (right, fitted to a spherical-harmonic approximation) isodensity surfaces calculated with ParaSurf™ using the default settings	6
Figure 2	2D-representation of a solvent-excluded surface	8
Figure 3	The solvent-accessible surface is obtained by rolling a spherical "solvent molecule"	8
Figure 4	2D-representation of a molecular surface with single-valued (A and B) and multiply valued (C and D) radial vectors from the center	9
Figure 5	2D-representation of the shrink-wrap algorithm. The algorithms scans along the vector from point a towards the center of the molecule until the electron density reaches the preset value (point b). The algorithm results in enclosures (marked yellow) for multi-valued radial vectors	9
Figure 6	Spherical-harmonic approximation of a shrink-wrap isodensity surface. Note the areas where the surface does not follow the indentations of the molecule	10
Figure 7	Schematic representation of the planes and hinge area used to determine the center for spherical-harmonic expansions	12
Figure 8	The four autocorrelation functions calculated using the AM1 Hamiltonian for trimethoprim	28
Figure 9	A sample <FRAGMENTS> input block	36
Figure 10	The fragments defined in the SDF input example	37
Figure 11	ParaSurf'12™ output for the phenyl fragment defined above	38
Figure 12	Surfaces calculated for the individual fragments, color coded according to the MEP in kcal mol ⁻¹ . The fragments (clockwise from the top right) are methoxy1, methoxy2, methoxy3, thymine, phenyl and methylene	39
Figure 13	The headers and titles, atomic coordinates and bond definitions from a VAMP .sdf file. The format follows the MDL definition [26]	41
Figure 14	ParaSurf™ output for trimethoprim, 1, using a spherical-harmonic surface	52
Figure 15	ParaSurf™ output for trimethoprim using a marching-cube surface	58
Figure 16	Shannon entropy section of the ParaSurf™ output for trimethoprim, 1, using a marching-cube isodensity surface.	59
Figure 17	Shannon entropy section of the ParaSurf™ output for trimethoprim, 1, using a marching-cube isodensity surface	60
Figure 18	Similarity output using autocorrelation functions. The lead molecule is captopril, which is defined in captopril_p.sdf	60
Figure 19	Sample grid file	70
Figure 20	Sample grid output file	71
Figure 21	Sample surface-integral model (SIM) file	73



8 REFERENCES

- [1] T. Clark, A. Alex, B. Beck, J. Chandrasekhar, P. Gedeck, A. H. C. Horn, M. Hutter, B. Martin, G. Rauhut, W. Sauer, T. Schindler, T. Steinke, VAMP 8.2, **Erlangen 2002**. Available from Accelrys Inc., San Diego, USA (<http://www.accelrys.com/products/datasheets/vamp.pdf>).
- [2] J. J. P. Stewart, MOPAC2000, **1999**, Fujitsu, Ltd, Tokyo, Japan. MOPAC 6.0 was once available as: J. J. P. Stewart, QCPE # 455, Quantum Chemistry Program Exchange, Bloomsburg, Indiana, 1990.
- [3] a) J. H. Van Drie. 'Shrink-wrap' surfaces: a new method for incorporating shape into pharmacophoric 3D database searching, *Journal of Chemical Information & Computer Sciences* **1997**, 37, 38-42; b) J. H. Van Drie, R. A. Nugent. Addressing the challenges of combinatorial chemistry: 3D databases, pharmacophore recognition and beyond, SAR and QSAR in Env. Res. **1998**, 9, 1-21; c) J. Erickson, D. J. Neidhart, J. H. Van Drie, D. J. Kempf, X. C. Wang, D. W. Norbeck, J. J. Plattner, J. W. Rittenhouse, M. Turon, N. Wideburg, et al. Design, activity, and 2.8 Å crystal structure of a C2 symmetric inhibitor complexed to HIV-1 protease, *Science* **1990**, 249, 527-533
- [4] W. Heiden, T. Goetze, J. Brickmann. Fast generation of molecular surfaces from 3D data fields with an enhanced "marching cube" algorithm, *J. Comput. Chem.* **1993**, 14, 246-250
- [5] D. W. Ritchie, G. J. L. Kemp. Fast computation, rotation, and comparison of low resolution spherical harmonic molecular surfaces, *J. Comput. Chem.* **1999**, 20, 383
- [6] *Chemical Applications of Atomic and Molecular Electrostatic Potentials. Reactivity, Structure, Scattering, and Energetics of Organic, Inorganic, and Biological Systems*, P. Politzer, D. G. Truhlar (Eds.), Plenum Press, New York, NY, **1981**.
- [7] P. Sjoberg, J. S. Murray, T. Brinck, P. Politzer. Average local ionisation energies on the molecular surfaces of aromatic systems as guides to chemical reactivity, *Can. J. Chem.* **1990**, 68, 1440-1443
- [8] B. Ehresmann, B. Martin, A. H. Horn, T. Clark. Local molecular properties and their use in predicting reactivity, *Journal of Molecular Modeling* **2003**, 9, 342-347. doi:10.1007/s00894-003-0153-x
- [9] B. Ehresmann, M. J. de Groot, A. Alex, T. Clark. New molecular descriptors based on local properties at the molecular surface and a boiling-point model derived from them, *J Chem Inf Comput Sci* **2004**, 44, 658-668. doi:10.1021/ci034215e
- [10] B. Ehresmann, M. J. de Groot, T. Clark. Surface-integral QSPR models: local energy properties, *J Chem Inf Model* **2005**, 45, 1053-1060. doi:10.1021/ci050025n
- [11] L. M. Loew, W. R. MacArthur. A molecular orbital study of monomeric metaphosphate. Density surfaces of frontier orbitals as a tool in assessing reactivity, *J. Am. Chem. Soc.* **1977**, 99, 1019-1025
- [12] B. S. Duncan, A. J. Olson. Approximation and visualization of large-scale motion of protein surfaces, *J Mol Graph* **1995**, 13, 250-257
- [13] J. H. Lin, T. Clark. An analytical, variable resolution, complete description of static molecules and their intermolecular binding properties, *J Chem Inf Model* **2005**, 45, 1010-1016. doi:10.1021/ci050059v
- [14] a) G. Rauhut, T. Clark. Multicenter Point Charge Model for High Quality Molecular Electrostatic Potentials from AM1 Calculations, *J. Comput. Chem.* **1993**, 14, 503-509; b) B. Beck, G. Rauhut, T. Clark. The Natural Atomic Orbital Point Charge Model for PM3: Multipole Moments and Molecular Electrostatic Potentials, *J. Comput. Chem.* **1994**, 15, 1064-1073
- [15] a) M. J. S. Dewar, W. Thiel. MNDO, *J. Am. Chem. Soc.* **1977**, 99, 4899-4907; 4907-4917; b) W. Thiel. MNDO, in *Encyclopedia of Computational Chemistry*, Vol. 1, P. v. R. Schleyer, N. L. Allinger, T. Clark, J. Gasteiger, P. A. Kollman, H. F. Schaefer, III, P. R. Schreiner (Eds.), Wiley, Chichester, **1998**, p. 1599
- [16] a) M. J. S. Dewar, E. G. Zoebisch, E. F. Healy, J. J. P. Stewart. Development and use of quantum mechanical molecular models. 76. AM1: a new general purpose quantum mechanical molecular model, *Journal of the American Chemical Society* **1985**, 107, 3902-3909. doi:10.1021/ja00299a024; b) A. J. Holder. AM1, in *Encyclopedia of Computational Chemistry*, Vol. 1, P. v. R. Schleyer, N. L. Allinger, T. Clark, J. Gasteiger, P. A. Kollman, H. F. Schaefer, III, P. R. Schreiner (Eds.), Wiley, Chichester, **1998**, pp. 8-11
- [17] a) J. J. P. Stewart. Optimization of parameters for semiempirical methods II. Applications, *Journal of Computational Chemistry* **1989**, 10, 221-264. doi:10.1002/jcc.540100209; b) J. J. Stewart. PM3, in *Encyclopedia of Computational Chemistry*, Vol. 1, P. v. R. Schleyer, N. L. Allinger, T. Clark, J. Gasteiger, P. A. Kollman, H. F. Schaefer, III, P. R. Schreiner (Eds.), Wiley, Chichester, **1998**, pp. 2080-2086



- [18] P. Winget, A. C. Horn, C. Selçuki, B. Martin, T. Clark. AM1* parameters for phosphorus, sulfur and chlorine, *Journal of Molecular Modeling* **2003**, 9, 408-414. doi:10.1007/s00894-003-0156-7
- [19] a) W. Thiel, A. A. Voityuk. Extension of the MNDO formalism to d orbitals: integral approximations and preliminary numerical results, *Theor. Chim. Acta* **1992**, 81, 391-404; b) W. Thiel, A. A. Voityuk. Extension of MNDO to d orbitals: parameters and results for the halogens, *Theor. Chim. Acta* **1996**, 93, 315-315; c) W. Thiel, A. A. Voityuk. Extension of MNDO to d orbitals: parameters and results for the silicon, *J. Mol. Struct.* **1994**, 313, 141-154; d) W. Thiel, A. A. Voityuk. Extension of MNDO to d Orbitals: Parameters and Results for the Second-Row Elements and for the Zinc Group, *The Journal of Physical Chemistry* **1996**, 100, 616-626. doi:10.1021/jp952148o; e) W. Thiel. MNDO/d, in *Encyclopedia of Computational Chemistry*, Vol. 3, P. v. R. Schleyer, N. L. Allinger, T. Clark, J. Gasteiger, P. A. Kollman, H. F. Schaefer, III, P. R. Schreiner (Eds.), Wiley, Chichester, **1998**, pp. 1604-1606
- [20] a) T. Clark. The local electron affinity for non-minimal basis sets, *Journal of Molecular Modeling* **2010**, 16, 1231-1238. doi:10.1007/s00894-009-0607-x; b) B. Martin, M. Gedeck, T. Clark. An Additive NDDO-Based Atomic Polarizability Model, *Int. J. Quant. Chem.* **2000**, 77, 473-497
- [21] A. H. C. Horn, J.-H. Lin, T. Clark. A Multipole Electrostatic Model for NDDO-based Semiempirical Molecular Orbital Methods, *Theor. Chem. Accts.* **2005**, 113, 159-168
- [22] G. Schürer, M. Gedeck, M. Gottschalk, T. Clark. Accurate Parametrized Variational Calculations of the Molecular Electronic Polarizability by NDDO-Based Methods, *J. Quant. Chem.* **1999**, 75, 17
- [23] a) D. Rinaldi, J.-L. Rivail. Calculation of molecular electronic polarizabilities. Comparison of different methods, *Theor. Chim. Acta* **1974**, 32, 243-251; b) J.-L. Rivail, D. Rinaldi. Variational calculation of multipole electric polarizabilities, *ComptesRendus, Serie B: Sciences Physiques* **1976**, 283-283; c) D. Rinaldi, J.-L. Rivail. Molecular polarizabilities and dielectric effect of the medium in the liquid state. Theoretical study of the water molecule and its dimers, *Theor. Chim. Acta* **1973**, 32, 57-57
- [24] R. J. Abraham, B. D. Hudson, M. W. Kermode, J. R. Mines. A general calculation of molecular solvation energies, *Journal of the Chemical Society, Faraday Transactions 1: Physical Chemistry in Condensed Phases* **1988**, 84, 1911-1917. doi:10.1039/f19888401911
- [25] J. G. Cramer, G. R. Famini, A. H. Lowrey. *Acc. Chem. Res.* **1993**, 26, 599-605
- [26] A. Y. Meyer. *Chem. Soc. Rev.* **1986**, 15, 449-475
- [27] M. Wagener, J. Sadowski, J. Gasteiger. Autocorrelation of Molecular Surface Properties for Modeling Corticosteroid Binding Globulin and Cytosolic A Receptor Activity by Neural Networks, *J. Am. Chem. Soc.* **1995**, 117, 7769-7775
- [28] CORINA, Molecular Networks GmbH, Erlangen. <http://www.molecular-networks.com/products/corina>
- [29] LogKOW dataset. <http://logkow.cisti.nrc.ca/logkow/>
- [30] M. Muehlbacher, A. E. Kerdawy, C. Kramer, B. Hudson, T. Clark. Conformation-Dependent QSPR Models: logPOW, *J Chem Inf Model* **2011**, 51, 2408-2416. doi:10.1021/ci200276v
- [31] C. E. Shannon, W. Weaver, *The Mathematical Theory of Communication*, University of Illinois Press, Chicago, **1949**.
- [32] a) R. Wang, X. Fang, Y. Lu, C. Y. Yang, S. Wang. The PDBbind database: methodologies and updates, *J Med Chem* **2005**, 48, 4111-4119. doi:10.1021/jm048957q; b) R. Wang, X. Fang, Y. Lu, S. Wang. The PDBbind database: collection of binding affinities for protein-ligand complexes with known three-dimensional structures, *J Med Chem* **2004**, 47, 2977-2980. doi:10.1021/jm030580l; c) PDBbind database. <http://www.pdbbind.org/>
- [33] M. Connelly. The molecular surface package, *J. Mol. Graph.* **1992**, 11, 139-141
- [34] L. Mavridis, B. D. Hudson, D. W. Ritchie. Toward high throughput 3D virtual screening using spherical harmonic surface representations, *J Chem Inf Model* **2007**, 47, 1787-1796. doi:10.1021/ci7001507
- [35] A. J. Jakobi, H. Mauser, T. Clark. ParaFrag--an approach for surface-based similarity comparison of molecular fragments, *Journal of Molecular Modeling* **2008**, 14, 547-558. doi:10.1007/s00894-008-0302-3
- [36] a) M. Hennemann, T. Clark. A QSPR-approach to the estimation of the pK(HB) of six-membered nitrogen-heterocycles using quantum mechanically derived descriptors, *Journal of Molecular Modeling* **2002**, 8, 95-101. doi:10.1007/s00894-002-0075-z; b) M. Hennemann, A. Friedl, M. Lobell, J. Keldenich, A. Hillisch, T. Clark, A. H. Goller. CypScore: Quantitative prediction of reactivity toward cytochromes P450 based on semiempirical molecular orbital theory, *ChemMedChem* **2009**, 4, 657-669. doi:10.1002/cmdc.200800384
- [37] A. Bondi. van der Waals Volumes and Radii, *J. Phys. Chem.* **1964**, 68, 441-451
- [38] http://www.mdl.com/solutions/white_papers/ctfile_formats.jsp



- [39] W. Thiel. The MNDOC method, a correlated version of the MNDO model, *J. Am. Chem. Soc.* **1981**, *103*, 1413-1420
- [40] R. J. Abraham, B. D. Hudson, M. W. Kermode, J. R. Mines. *Faraday Trans. I* **1988**, *84*, 1911-1917
- [41] T. Clark, K. G. Byler, M. J. de Groot. Biological Communication via Molecular Surfaces, in *Molecular Interactions - Bringing Chemistry to Life (Proceedings of the International Beilstein Workshop, Bozen, Italy, May 15-19, 2006)*, Logos Verlag, Bozen, Italy, **2008**. (Available at <http://www.beilstein-institut.de/bozen2006/proceedings/Clark/Clark.pdf>)